



## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification <sup>6</sup> :</b> <b>C12N 15/12, C07K 14/47, 14/495, C12N 15/62, A61K 38/17, C07K 16/18, C12Q 1/68</b>	<b>A2</b>	<b>(11) International Publication Number:</b> <b>WO 99/33979</b> <b>(43) International Publication Date:</b> 8 July 1999 (08.07.99)
<b>(21) International Application Number:</b> PCT/US98/27008 <b>(22) International Filing Date:</b> 18 December 1998 (18.12.98)  <b>(30) Priority Data:</b> 60/068,958 30 December 1997 (30.12.97) US 60/101,603 24 September 1998 (24.09.98) US 60/102,540 30 September 1998 (30.09.98) US  <b>(71) Applicant:</b> CHIRON CORPORATION [US/US]; 4560 Horton Street - R440, Emeryville, CA 94608 (US).  <b>(72) Inventors:</b> LIN, Haishan; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94622-8097 (US). CAO, Li; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94622-8097 (US).  <b>(74) Agents:</b> POTTER, Jane, E., R. et al.; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US).		<b>(81) Designated States:</b> AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).  <b>Published</b> <i>Without international search report and to be republished upon receipt of that report.</i>
<b>(54) Title:</b> BONE MARROW SECRETED PROTEINS AND POLYNUCLEOTIDES  <b>(57) Abstract</b>  Novel polynucleotides and secreted proteins encoded thereby are disclosed. The proteins can be used as therapeutics, for example, to stimulate blood cell generation in patients receiving cancer chemotherapy, to treat bone marrow transplantation patients, and to heal fractured bones. Polynucleotides of the invention can be used therapeutically, to provide proteins of the invention. Polynucleotides of the invention can also be used diagnostically, such as on polynucleotide arrays, to detect differential gene expression in diseased tissue compared with gene expression in normal tissue.		

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Larvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

## **BONE MARROW SECRETED PROTEINS AND POLYNUCLEOTIDES**

### **5     TECHNICAL AREA OF THE INVENTION**

This invention relates to proteins secreted from bone marrow and to polynucleotides encoding the secreted proteins. The invention also relates to therapeutic and diagnostic utilities for the polynucleotides and proteins.

### **10    BACKGROUND OF THE INVENTION**

Bone marrow stromal cells secrete a variety of protein factors required for the formation of blood and bone cells and for other physiological processes. Known regulatory factors involved in hematopoiesis and/or bone development include SCF, IL-3, IL-6, GM-CSF, M-CSF, EPO, TPO, bone morphogenic proteins, erythroid potentiating factor, and TGF-  
15     $\beta$ . However, it is believed that additional secreted protein factors which control hematopoiesis and bone morphogenesis remain to be identified.

### **SUMMARY OF THE INVENTION**

It is an object of the invention to provide proteins secreted from bone marrow  
20    stromal cells and polynucleotides encoding the secreted proteins. These and other objects of the invention are provided by one or more of the embodiments described below.

One embodiment of the invention is an isolated and purified protein comprising an amino acid sequence which is at least 85% identical to an amino acid sequence selected  
25    from the group consisting of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, and 44. Percent identity is determined using a Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 1.

Another embodiment of the invention is an isolated and purified protein comprising an  
30    amino acid sequence selected from the group consisting of at least 95 contiguous amino

acids of SEQ ID NO:2, at least 101 contiguous amino acids of SEQ ID NO:4, at least 14  
contiguous amino acids selected from amino acids 1-312 of SEQ ID NO:6, at least 179  
contiguous amino acids of SEQ ID NO:6, at least 75 contiguous amino acids of SEQ ID  
NO:6, at least 179 contiguous amino acids of SEQ ID NO:8, at least 136 contiguous  
5 amino acids of SEQ ID NO:8, at least 17 contiguous amino acids selected from amino  
acids 1-287 of SEQ ID NO:8, at least 82 contiguous amino acids of SEQ ID NO:10, at  
least 31 contiguous amino acids selected from amino acids 1-238 of SEQ ID NO:10, at  
least 96 contiguous amino acids of SEQ ID NO:12, at least 27 contiguous amino acids  
selected from amino acids 250-383 of SEQ ID NO:12, at least 6 contiguous amino acids  
10 selected from amino acids 1-184 of SEQ ID NO:12, at least 8 contiguous amino acids  
selected from amino acids 268-364 of SEQ ID NO:12, at least 104 contiguous amino  
acids of SEQ ID NO:14, at least 75 contiguous amino acids of SEQ ID NO:14, at least 17  
contiguous amino acids selected from amino acids 1-150 of SEQ ID NO:14, at least 6  
contiguous amino acids selected from amino acids 204-261 of SEQ ID NO:14, at least 6  
15 contiguous amino acids selected from amino acids 1-111 of SEQ ID NO:14, at least 8  
contiguous amino acids of SEQ ID NO:16, at least 46 contiguous amino acids of SEQ ID  
NO:18, at least 39 contiguous amino acids selected from amino acids 13-232 of SEQ ID  
NO:18, at least 6 contiguous amino acids of SEQ ID NO:20, at least 7 contiguous amino  
acids of SEQ ID NO:22, at least 7 contiguous amino acids of SEQ ID NO:24, at least 11  
20 contiguous amino acids of SEQ ID NO:26, at least 257 contiguous amino acids of SEQ  
ID NO:28, at least 6 contiguous amino acids selected from amino acids 1-31 of SEQ ID  
NO:28, at least 6 contiguous amino acids of SEQ ID NO:30, at least 117 contiguous  
amino acids of SEQ ID NO:32, at least 6 contiguous amino acids selected from amino  
acids 1-65 of SEQ ID NO:32, at least 6 contiguous amino acids of SEQ ID NO:34, at  
25 least 14 contiguous amino acids of SEQ ID NO:36, at least 19 contiguous amino acids of  
SEQ ID NO:38, at least 8 contiguous amino acids of SEQ ID NO:40, at least 7  
contiguous amino acids of SEQ ID NO:42, and at least 10 contiguous amino acids of  
SEQ ID NO:44.

Still another embodiment of the invention is a fusion protein comprising two  
30 protein segments joined together with a peptide bond. The first protein segment consists

of an amino acid sequence selected from the group consisting of at least 95 contiguous amino acids of SEQ ID NO:2, at least 101 contiguous amino acids of SEQ ID NO:4, at least 14 contiguous amino acids selected from amino acids 1-312 of SEQ ID NO:6, at least 179 contiguous amino acids of SEQ ID NO:6, at least 75 contiguous amino acids of SEQ ID NO:6, at least 179 contiguous amino acids of SEQ ID NO:8, at least 136 contiguous amino acids of SEQ ID NO:8, at least 17 contiguous amino acids selected from amino acids 1-287 of SEQ ID NO:8, at least 82 contiguous amino acids of SEQ ID NO:10, at least 31 contiguous amino acids selected from amino acids 1-238 of SEQ ID NO:10, at least 96 contiguous amino acids of SEQ ID NO:12, at least 27 contiguous amino acids selected from amino acids 250-383 of SEQ ID NO:12, at least 6 contiguous amino acids selected from amino acids 1-184 of SEQ ID NO:12, at least 8 contiguous amino acids selected from amino acids 268-364 of SEQ ID NO:12, at least 104 contiguous amino acids of SEQ ID NO:14, at least 75 contiguous amino acids of SEQ ID NO:14, at least 17 contiguous amino acids selected from amino acids 1-150 of SEQ ID NO:14, at least 6 contiguous amino acids selected from amino acids 204-261 of SEQ ID NO:14, at least 6 contiguous amino acids selected from amino acids 1-111 of SEQ ID NO:14, at least 8 contiguous amino acids of SEQ ID NO:16, at least 46 contiguous amino acids of SEQ ID NO:18, at least 39 contiguous amino acids selected from amino acids 13-232 of SEQ ID NO:18, at least 6 contiguous amino acids of SEQ ID NO:20, at least 7 contiguous amino acids of SEQ ID NO:22, at least 7 contiguous amino acids of SEQ ID NO:24, at least 11 contiguous amino acids of SEQ ID NO:26, at least 257 contiguous amino acids of SEQ ID NO:28, at least 6 contiguous amino acids selected from amino acids 1-31 of SEQ ID NO:28, at least 6 contiguous amino acids of SEQ ID NO:30, at least 117 contiguous amino acids of SEQ ID NO:32, at least 6 contiguous amino acids selected from amino acids 1-65 of SEQ ID NO:32, at least 6 contiguous amino acids of SEQ ID NO:34, at least 14 contiguous amino acids of SEQ ID NO:36, at least 19 contiguous amino acids of SEQ ID NO:38, at least 8 contiguous amino acids of SEQ ID NO:40, at least 7 contiguous amino acids of SEQ ID NO:42, and at least 10 contiguous amino acids of SEQ ID NO:44.

Even another embodiment of the invention is a preparation of antibodies which

specifically binds to a protein comprising an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, and 44.

5 Still another embodiment of the invention is an isolated and purified subgenomic polynucleotide which encodes a protein comprising an amino acid sequence which is at least 85% identical to an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, and 44. Percent identity is determined using a Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of  
10 1.

A further embodiment of the invention is an isolated and purified subgenomic polynucleotide comprising a nucleotide sequence which is at least 85% identical to a nucleotide sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 31, 33, 35, 37, 39, 41, 43, 45, and the complements thereof.  
15 Percent identity is determined using a Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 1.

Another embodiment of the invention is an isolated and purified subgenomic polynucleotide which encodes an amino acid sequence selected from the group  
20 consisting of at least 95 contiguous amino acids of SEQ ID NO:2, at least 101 contiguous amino acids of SEQ ID NO:4, at least 14 contiguous amino acids selected from amino acids 1-312 of SEQ ID NO:6, at least 179 contiguous amino acids of SEQ ID NO:6, at least 75 contiguous amino acids of SEQ ID NO:6, at least 179 contiguous amino acids of SEQ ID NO:8, at least 136 contiguous amino acids of SEQ ID NO:8, at least 17  
25 contiguous amino acids selected from amino acids 1-287 of SEQ ID NO:8, at least 82 contiguous amino acids of SEQ ID NO:10, at least 31 contiguous amino acids selected from amino acids 1-238 of SEQ ID NO:10, at least 96 contiguous amino acids of SEQ ID NO:12, at least 27 contiguous amino acids selected from amino acids 250-383 of SEQ ID NO:12, at least 6 contiguous amino acids selected from amino acids 1-184 of SEQ ID  
30 NO:12, at least 8 contiguous amino acids selected from amino acids 268-364 of SEQ ID

NO:12, at least 104 contiguous amino acids of SEQ ID NO:14, at least 75 contiguous amino acids of SEQ ID NO:14, at least 17 contiguous amino acids selected from amino acids 1-150 of SEQ ID NO:14, at least 6 contiguous amino acids selected from amino acids 204-261 of SEQ ID NO:14, at least 6 contiguous amino acids selected from amino acids 1-111 of SEQ ID NO:14, at least 8 contiguous amino acids of SEQ ID NO:16, at least 46 contiguous amino acids of SEQ ID NO:18, at least 39 contiguous amino acids selected from amino acids 13-232 of SEQ ID NO:18, at least 6 contiguous amino acids of SEQ ID NO:20, at least 7 contiguous amino acids of SEQ ID NO:22, at least 7 contiguous amino acids of SEQ ID NO:24, at least 11 contiguous amino acids of SEQ ID NO:26, at least 257 contiguous amino acids of SEQ ID NO:28, at least 6 contiguous amino acids selected from amino acids 1-31 of SEQ ID NO:28, at least 6 contiguous amino acids of SEQ ID NO:30, at least 117 contiguous amino acids of SEQ ID NO:32, at least 6 contiguous amino acids selected from amino acids 1-65 of SEQ ID NO:32, at least 6 contiguous amino acids of SEQ ID NO:34, at least 14 contiguous amino acids of SEQ ID NO:36, at least 19 contiguous amino acids of SEQ ID NO:38, at least 8 contiguous amino acids of SEQ ID NO:40, at least 7 contiguous amino acids of SEQ ID NO:42, and at least 10 contiguous amino acids of SEQ ID NO:44.

Still another embodiment of the invention is an isolated and purified subgenomic polynucleotide comprising a polynucleotide segment which hybridizes to a nucleotide sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 31, 33, 35, 37, 39, 41, and 43, and the complements thereof after washing with 0.2X SSC at 65 °C, wherein the polynucleotide segment encodes a protein having an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, and 44.

Even another embodiment of the invention is an isolated and purified subgenomic polynucleotide comprising a nucleotide sequence selected from the group consisting of at least 499 contiguous nucleotides of SEQ ID NO:1, at least 1141 contiguous nucleotides of SEQ ID NO:1, at least 475 contiguous nucleotides of SEQ ID NO:3, at least 313 contiguous nucleotides selected from nucleotides 1-1001 of SEQ ID NO:3, at least 751 contiguous nucleotides of SEQ ID NO:5, at least 538 contiguous nucleotides of SEQ ID

NO:5, at least 11 contiguous nucleotides selected from nucleotides 1-946 of SEQ ID  
NO:5, at least 13 contiguous nucleotides selected from nucleotides 1-1039 of SEQ ID  
NO:5, at least 651 contiguous nucleotides of SEQ ID NO:7, at least 522 contiguous  
nucleotides of SEQ ID NO:7, at least 11 contiguous nucleotides selected from  
5 nucleotides 1-913 of SEQ ID NO:7, at least 484 contiguous nucleotides of SEQ ID  
NO:9, at least 317 contiguous nucleotides of SEQ ID NO:9, at least 11 contiguous  
nucleotides selected from nucleotides 1-216 of SEQ ID NO:9, at least 11 contiguous  
nucleotides selected from nucleotides 379-812 of SEQ ID NO:9, at least 183 contiguous  
nucleotides selected from nucleotides 1-984 of SEQ ID NO:9, at least 594 contiguous  
10 nucleotides of SEQ ID NO:11, at least 289 contiguous nucleotides of SEQ ID NO:11, at  
least 11 contiguous nucleotides selected from nucleotides 1-585 of SEQ ID NO:11, at  
least 11 contiguous nucleotides selected from nucleotides 853-1120 of SEQ ID NO:11, at  
least 592 contiguous nucleotides of SEQ ID NO:13, at least 275 contiguous nucleotides  
of SEQ ID NO:13, at least 11 contiguous nucleotides selected from nucleotides 1-294 of  
15 SEQ ID NO:13, at least 537 contiguous nucleotides of SEQ ID NO:15, at least 294  
contiguous nucleotides selected from nucleotides 1-1889 of SEQ ID NO:15, at least 171  
contiguous nucleotides selected from nucleotides 318-1766 of SEQ ID NO:15, at least 11  
contiguous nucleotides selected from nucleotides 1-42 of SEQ ID NO:15, at least 11  
contiguous nucleotides selected from nucleotides 478-908 of SEQ ID NO:15, at least 11  
20 contiguous nucleotides selected from nucleotides 1059-1078 of SEQ ID NO:15, at least  
205 contiguous nucleotides of SEQ ID NO:17, at least 440 contiguous nucleotides of  
SEQ ID NO:19, at least 451 contiguous nucleotides of SEQ ID NO:21, at least 11  
contiguous nucleotides selected from nucleotides 1-121 of SEQ ID NO:21, at least 11  
contiguous nucleotides selected from nucleotides 474-592 of SEQ ID NO:21, at least 351  
25 contiguous nucleotides of SEQ ID NO:23, at least 21 contiguous nucleotides selected  
from nucleotides 1-1943 of SEQ ID NO:23, at least 11 contiguous nucleotides selected  
from 1-612 of SEQ ID NO:23, at least 11 contiguous nucleotides selected from  
nucleotides 611-719 of SEQ ID NO:23, at least 11 contiguous nucleotides selected from  
nucleotides 713-830 of SEQ ID NO:23, at least 11 contiguous nucleotides selected from  
30 nucleotides 830-1933 of SEQ ID NO:23, at least 492 nucleotides of SEQ ID NO:25, at



least 11 contiguous nucleotides selected from nucleotides 758-847 of SEQ ID NO:25, at least 1024 contiguous nucleotides of SEQ ID NO:27, at least 347 contiguous nucleotides of SEQ ID NO: 29, at least 11 contiguous nucleotides selected from nucleotides 548-601 of SEQ ID NO:29, at least 394 contiguous nucleotides of SEQ ID NO: 31, at least 11  
5 contiguous nucleotides selected from nucleotides 1-361 of SEQ ID NO:31, at least 11 contiguous nucleotides selected from nucleotides 1083-1102 of SEQ ID NO:31, at least 492 contiguous nucleotides of SEQ ID NO:33, at least 510 contiguous nucleotides of SEQ ID NO:35, at least 11 contiguous nucleotides selected from nucleotides 1-502 or 505-631 of SEQ ID NO:35, at least 392 contiguous nucleotides of SEQ ID NO:37, at  
10 least 11 contiguous nucleotides selected from nucleotides 1-502 of SEQ ID NO:37, at least 11 contiguous nucleotides selected from nucleotides 505-631 of SEQ ID NO:37, at least 559 contiguous nucleotides of SEQ ID NO:39, at least 11 contiguous nucleotides selected from nucleotides 1-92 of SEQ ID NO:39, at least 254 contiguous nucleotides of SEQ ID NO:41, at least 11 contiguous nucleotides selected from nucleotides 1-34 of  
15 SEQ ID NO:41 at least 11 contiguous nucleotides selected from nucleotides 55-110 of SEQ ID NO:41, at least 103 contiguous nucleotides of SEQ ID NO:43, at least 11 contiguous nucleotides selected from nucleotides 1-280 of SEQ ID NO:43, at least 11 contiguous nucleotides selected from nucleotides 270-319 of SEQ ID NO:43, at least 11 contiguous nucleotides selected from nucleotides 378-423 of SEQ ID NO:43, at least 11  
20 contiguous nucleotides selected from nucleotides 414-492 of SEQ ID NO:43, at least 11 contiguous nucleotides selected from nucleotides 532-570 of SEQ ID NO:43, at least 11 contiguous nucleotides selected from nucleotides 1086-1152 of SEQ ID NO:43, and the complements thereof.

A further embodiment of the invention is a construct comprising isolated and  
25 purified subgenomic polynucleotides of the invention.

Another embodiment of the invention is a host cell comprising a construct of the invention.

Yet another embodiment of the invention is a process for producing a protein. A culture of a host cell comprising a construct of the invention is grown in a suitable  
30 culture medium. The protein secreted from the host cell is purified.

Another embodiment of the invention is a polynucleotide array comprising at least one single-stranded polynucleotide which comprises at least 12 contiguous nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, and the complements thereof.

Even another embodiment of the invention is a method of detecting differential gene expression between two biological samples. A first biological sample comprising single-stranded polynucleotide molecules with a first polynucleotide array comprising at least one single-stranded polynucleotide which comprises at least 12 contiguous nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, 43, and the complements thereof. A second biological sample comprising single-stranded polynucleotide molecules is contacted with a second polynucleotide array. The first and second polynucleotide arrays comprise identical single-stranded polynucleotides. A first and second pattern of double-stranded polynucleotides bound to the first and second polynucleotide arrays are detected. A difference between the first and second patterns indicates a gene which is differentially expressed between the first and second biological samples.

Methods are also provided for preventing, treating, or ameliorating a medical condition associated with hematopoiesis or bone marrow morphogenesis, which comprises administering to a mammalian subject a therapeutically effective amount of a composition comprising a protein of the present invention and a pharmaceutically acceptable carrier.

Proteins encoded by polynucleotides of the present invention have potential uses in stimulating blood cell generation in patient receiving cancer chemotherapy, for bone marrow transplantation patient, and for healing fractured bones.

#### **DETAILED DESCRIPTION OF THE INVENTION**

Secreted proteins include proteins which, when expressed in a suitable host cell, are transported across or through a membrane, including transport as a result of signal

sequences. Secreted proteins include proteins which are secreted wholly (*e.g.*, soluble proteins) or partially (*e.g.*, receptors) from the cell in which they are expressed. Secreted proteins also include proteins which are transported across the membrane of the endoplasmic reticulum.

5 Polynucleotides of the invention which encode secreted proteins were isolated from a cDNA library derived from human bone marrow stromal cells. Subgenomic polynucleotides of the invention contain less than a whole chromosome and can be single- or double-stranded. Preferably, the polynucleotides are intron-free. Subgenomic polynucleotides of the invention can comprise all or a portion of a nucleotide sequence  
10 disclosed in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, or 43, as explained in detail below. The complements of these nucleotide sequences are contiguous nucleotide sequences which form Watson-Crick base pairs with a contiguous nucleotide sequence as shown in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, or 43. These complementary sequences are  
15 also subgenomic polynucleotides and can be used, *inter alia*, to provide antisense oligonucleotides.

Degenerate nucleotide sequences encoding amino acid sequences of proteins of the invention, as well as homologous nucleotide sequences which are at least 65%, 75%, 85%, 90%, 95%, 98%, or 99% identical to the nucleotide sequences shown in NOS:1, 3,  
20 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, and 43, are also subgenomic polynucleotides of the invention. Percent identity is determined using computer programs which employ the Smith-Waterman homology search algorithm, for example as implemented in the MPSRCH program (Oxford Molecular), using an affine gap search with the following parameters: a gap open penalty of 12 and a gap extension  
25 penalty of 1. The Smith-Waterman algorithm is taught in Smith and Waterman, *Adv. Appl. Math.* (1981) 2:482-489.

Typically, homologous sequences can be confirmed by hybridization under stringent conditions, as is known in the art. For example, using the following wash conditions-2X SSC (0.3 M NaCl, 0.03 M sodium citrate, pH 7.0), 0.1% SDS, room  
30 temperature twice, 30 minutes each; then 2X SSC, 0.1% SDS, 50 °C once, 30 minutes;

then 2X SSC, room temperature twice, 10 minutes each-homologous sequences can be identified which contain at most about 25-30% basepair mismatches. More preferably, homologous nucleic acid strands contain 15-25% basepair mismatches, even more preferably 5-15% basepair mismatches.

5           Species homologs of subgenomic polynucleotides of the invention can also be identified by making suitable probes or primers and screening cDNA expression libraries from other species, such as mice, monkeys, yeast, or bacteria, as well as human cDNA expression libraries. It is well known that the  $T_m$  of a double-stranded DNA decreases by 1-1.5 °C with every 1% decrease in homology (Bonner *et al.*, *J. Mol. Biol.* 81, 123  
10 (1973). Homologous subgenomic polynucleotide species can therefore be identified, for example, by hybridizing a putative homologous polynucleotide with a polynucleotide having a nucleotide sequence disclosed in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, or 43 to form a test hybrid, comparing the melting temperature of the test hybrid with the melting temperature of a hybrid  
15 comprising a polynucleotide having one of the disclosed nucleotide sequences and a polynucleotide which is perfectly complementary to that sequence, and calculating the number or percent of basepair mismatches within the test hybrid.

Nucleotide sequences which hybridize to the coding sequences shown in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, or 43 or  
20 their complements following stringent hybridization and/or wash conditions are also subgenomic polynucleotides of the invention. Stringent wash conditions are well known and understood in the art and are disclosed, for example, in Sambrook *et al.*, MOLECULAR CLONING: A LABORATORY MANUAL, 2d ed., 1989, at pages 9.50-9.51.

Typically, for stringent hybridization conditions a combination of temperature  
25 and salt concentration should be chosen that is approximately 12-20 °C below the calculated  $T_m$  of the hybrid under study. The  $T_m$  of a hybrid between a nucleotide sequence shown in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, or 43 and a polynucleotide sequence which is 65%, 75%, 85%, 90%, 95%, 96%, 97%, 98%, or 99% identical to that sequence can be calculated, for example,  
30 using the equation of Bolton and McCarthy, *Proc. Natl. Acad. Sci. U.S.A.* 48, 1390

(1962):

$$T_m = 81.5^{\circ}\text{C} - 16.6(\log_{10}[\text{Na}^+]) + 0.41(\%G + C) - 0.63(\%\text{formamide}) - 600/l,$$

where  $l$  = the length of the hybrid in basepairs.

Stringent wash conditions include, for example, 4X SSC at 65 °C, or 50% formamide, 4X  
5 SSC at 42 °C, or 0.5X SSC, 0.1% SDS at 65 °C. Highly stringent wash conditions  
include, for example, 0.2X SSC at 65 °C.

Subgenomic polynucleotides can be isolated and purified free from other  
nucleotide sequences using standard nucleic acid purification techniques. For example,  
restriction enzymes and probes can be used to isolate polynucleotide fragments which  
10 comprise nucleotide sequences of the invention. Isolated and purified subgenomic  
polynucleotides are in preparations which are free or at least 90% free of other  
molecules.

Complementary DNA (cDNA) molecules with coding sequences corresponding  
to SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41,  
15 or 43 are also subgenomic polynucleotides of the invention. cDNA molecules of the  
invention can be made with standard molecular biology techniques, using human mRNA  
as a template. cDNA molecules can thereafter be replicated using molecular biology  
techniques known in the art and disclosed in manuals such as Sambrook *et al.*, 1989. An  
amplification technique, such as the polymerase chain reaction (PCR), can be used to  
20 obtain additional copies of subgenomic polynucleotides of the invention, using either  
human genomic DNA or cDNA as a template.

Alternatively, synthetic chemistry techniques can be used to synthesize  
subgenomic polynucleotide molecules of the invention. The degeneracy of the genetic  
code allows alternate nucleotide sequences to be synthesized which will encode a protein  
25 having an amino acid sequence shown in SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20,  
22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, or 44 or a biologically active variant of one of  
those sequences. All such nucleotide sequences are within the scope of the present  
invention.

The invention also provides polynucleotide probes which can be used, for  
30 example, in hybridization protocols such as Northern or Southern blotting or *in situ*

hybridizations. Polynucleotide probes of the invention comprise at least 12, 13, 14, 15, 16, 17, 18, 19, 20, 30, or 40 or more contiguous nucleotides selected from SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, or 43.

Polynucleotide probes of the invention can comprise a detectable label, such as a  
5 radioisotopic, fluorescent, enzymatic, or chemiluminescent label.

Subgenomic polynucleotides of the invention can be used as primers to obtain additional copies of the polynucleotides. Subgenomic polynucleotides of the invention can also be used to express mRNA, protein, polypeptides, antibodies, or fusion proteins of the invention and to generate antisense oligonucleotides and ribozymes.

10 Isolated polynucleotides of the invention can be present in constructs, such as DNA or RNA constructs. They can be operably linked to a promoter or other expression control sequence in order to produce proteins of the invention recombinantly. Many suitable expression control sequences, such as the pMT2 or pED expression vectors disclosed in Kaufman *et al.*, *Nucleic Acids Res.* 19, 4485-4490 (1991), are well known in  
15 the art. General methods of expressing recombinant proteins are also well known (*see, e.g.*, Kaufman, *METHODS IN ENZYMOLOGY* 185, 537-566, 1990). An isolated polynucleotide and a promoter or an expression control sequence are operably linked when the isolated polynucleotide and the promoter or expression control sequence are situated within a construct or cell in such a way that the protein is expressed by a host  
20 cell which has been transformed or transfected with the polynucleotide and the promoter or expression control sequence.

For example, a construct of the invention can comprise a promoter which is functional in a particular type of host cell. The skilled artisan can readily select an appropriate promoter from the large number of cell type-specific promoters known and  
25 used in the art. The polynucleotide is located downstream from the promoter. Constructs of the invention can also contain a transcription terminator which is functional in the host cell. Transcription of the polynucleotide segment initiates at the promoter. A construct can be linear or circular and can contain sequences, if desired, for autonomous replication.

30 A variety of host cells are available for use in bacterial, yeast, insect, and human

expression systems and can be used to propagate or to express polynucleotides of the invention. Constructs comprising the polynucleotides can be introduced into host cells using any technique known in the art. These techniques include transferrin-polycation-mediated DNA transfer, transfection with naked or encapsulated nucleic acids, liposome-mediated cellular fusion, intracellular transportation of DNA-coated latex beads, 5 protoplast fusion, viral infection, electroporation, and calcium phosphate-mediated transfection.

Polynucleotides of the invention can be propagated in constructs and cell lines using techniques well known in the art. Polynucleotides can be on linear or circular 10 molecules. They can be on autonomously replicating molecules or on molecules without replication sequences. They can be regulated by their own or by other regulatory sequences, as are known in the art.

Bacterial systems for expressing polynucleotides of the invention include those described in Chang *et al.*, *Nature* (1978) 275: 615, Goeddel *et al.*, *Nature* (1979) 281: 15 544, Goeddel *et al.*, *Nucleic Acids Res.* (1980) 8: 4057, EP 36,776, U.S. 4,551,433, deBoer *et al.*, *Proc. Natl. Acad. Sci. USA* (1983) 80: 21-25, and Siebenlist *et al.*, *Cell* (1980) 20: 269.

Expression systems in yeast include those described in Hinnen *et al.*, *Proc. Natl. Acad. Sci. USA* (1978) 75: 1929; Ito *et al.*, *J. Bacteriol.* (1983) 153: 163; Kurtz *et al.*, 20 *Mol. Cell. Biol.* (1986) 6: 142; Kunze *et al.*, *J. Basic Microbiol.* (1985) 25: 141; Gleeson *et al.*, *J. Gen. Microbiol.* (1986) 132: 3459, Roggenkamp *et al.*, *Mol. Gen. Genet.* (1986) 202 :302) Das *et al.*, *J. Bacteriol.* (1984) 158: 1165; De Louvencourt *et al.*, *J. Bacteriol.* (1983) 154: 737, Van den Berg *et al.*, *Bio/Technology* (1990) 8: 135; Kunze *et al.*, *J. Basic Microbiol.* (1985) 25: 141; Cregg *et al.*, *Mol. Cell. Biol.* (1985) 5: 3376, U.S. 25 4,837,148, US 4,929,555; Beach and Nurse, *Nature* (1981) 300: 706; Davidow *et al.*, *Curr. Genet.* (1985) 10: 380, Gaillardin *et al.*, *Curr. Genet.* (1985) 10: 49, Ballance *et al.*, *Biochem. Biophys. Res. Commun.* (1983) 112: 284-289; Tilburn *et al.*, *Gene* (1983) 26: 205-221, Yelton *et al.*, *Proc. Natl. Acad. Sci. USA* (1984) 81: 1470-1474, Kelly and Hynes, *EMBO J.* (1985) 4: 475479; EP 244,234, and WO 91/00357.

30 Expression of polynucleotides of the invention in insects can be carried out as

described in U.S. 4,745,051, Friesen *et al.* (1986) "The Regulation of Baculovirus Gene Expression" in: THE MOLECULAR BIOLOGY OF BACULOVIRUSES (W. Doerfler, ed.), EP 127,839, EP 155,476, and Vlak *et al.*, *J. Gen. Virol.* (1988) 69: 765-776, Miller *et al.*, *Ann. Rev. Microbiol.* (1988) 42: 177, Carbonell *et al.*, *Gene* (1988) 73: 409, Maeda *et al.*, *Nature* (1985) 315: 592-594, Lebacqz-Verheyden *et al.*, *Mol. Cell. Biol.* (1988) 8: 3129; Smith *et al.*, *Proc. Natl. Acad. Sci. USA* (1985) 82: 8404, Miyajima *et al.*, *Gene* (1987) 58: 273; and Martin *et al.*, *DNA* (1988) 7:99. Numerous baculoviral strains and variants and corresponding permissive insect host cells from hosts are described in Luckow *et al.*, *Bio/Technology* (1988) 6: 47-55, Miller *et al.*, in GENETIC ENGINEERING (Setlow, J.K. *et al.* eds.), Vol. 8 (Plenum Publishing, 1986), pp. 277-279, and Maeda *et al.*, *Nature*, (1985) 315: 592-594.

Mammalian expression of polynucleotides can be achieved as described in Dijkema *et al.*, *EMBO J.* (1985) 4: 761, Gorman *et al.*, *Proc. Natl. Acad. Sci. USA* (1982b) 79: 6777, Boshart *et al.*, *Cell* (1985) 41: 521 and U.S. 4,399,216. Other features of mammalian expression can be facilitated as described in Ham and Wallace, *Meth. Enz.* (1979) 58: 44, Barnes and Sato, *Anal. Biochem.* (1980) 102: 255, U.S. 4,767,704, US 4,657,866, US 4,927,762, US 4,560,655, WO 90/103430, WO 87/00195, and U.S. RE 30,985.

Polynucleotides of the invention can also be used in gene delivery vehicles, for the purpose of delivering an mRNA or oligonucleotide (either with the sequence of a native mRNA or its complement), full-length protein, fusion protein, polypeptide, or ribozyme, or single-chain antibody, into a cell, preferably a eukaryotic cell. According to the present invention, a gene delivery vehicle can be, for example, naked plasmid DNA, a viral expression vector comprising a polynucleotide of the invention, or a polynucleotide of the invention in conjunction with a liposome or a condensing agent.

In one embodiment of the invention, the gene delivery vehicle comprises a promoter and one of the polynucleotides disclosed herein. Preferred promoters are tissue-specific promoters and promoters which are activated by cellular proliferation, such as the thymidine kinase and thymidylate synthase promoters. Other preferred promoters include promoters which are activatable by infection with a virus, such as the



$\alpha$ - and  $\beta$ -interferon promoters, and promoters which are activatable by a hormone, such as estrogen. Other promoters which can be used include the Moloney virus LTR, the CMV promoter, and the mouse albumin promoter.

A gene delivery vehicle can comprise viral sequences such as a viral origin of replication or packaging signal. These viral sequences can be selected from viruses such as astrovirus, coronavirus, orthomyxovirus, papovavirus, paramyxovirus, parvovirus, picornavirus, poxvirus, retrovirus, togavirus or adenovirus. In a preferred embodiment, the gene delivery vehicle is a recombinant retroviral vector. Recombinant retroviruses and various uses thereof have been described in numerous references including, for example, Mann *et al.*, *Cell* 33:153, 1983, Cane and Mulligan, *Proc. Nat'l. Acad. Sci. USA* 81:6349, 1984, Miller *et al.*, *Human Gene Therapy* 1:5-14, 1990, U.S. Patent Nos. 4,405,712, 4,861,719, and 4,980,289, and PCT Application Nos. WO 89/02,468, WO 89/05,349, and WO 90/02,806. Numerous retroviral gene delivery vehicles can be utilized in the present invention, including for example those described in EP 0,415,731; WO 90/07936; WO 94/03622; WO 93/25698; WO 93/25234; U.S. Patent No. 5,219,740; WO 9311230; WO 9310218; Vile and Hart, *Cancer Res.* 53:3860-3864, 1993; Vile and Hart, *Cancer Res.* 53:962-967, 1993; Ram *et al.*, *Cancer Res.* 53:83-88, 1993; Takamiya *et al.*, *J. Neurosci. Res.* 33:493-503, 1992; Baba *et al.*, *J. Neurosurg.* 79:729-735, 1993 (U.S. Patent No. 4,777,127, GB 2,200,651, EP 0,345,242 and WO91/02805).

Particularly preferred retroviruses are derived from retroviruses which include avian leukosis virus (ATCC Nos. VR-535 and VR-247), bovine leukemia virus (VR-1315), murine leukemia virus (MLV), mink-cell focus-inducing virus (Koch *et al.*, *J. Vir.* 49:828, 1984; and Oliff *et al.*, *J. Vir.* 48:542, 1983), murine sarcoma virus (ATCC Nos. VR-844, 45010 and 45016), reticuloendotheliosis virus (ATCC Nos VR-994, VR-770 and 45011), Rous sarcoma virus, Mason-Pfizer monkey virus, baboon endogenous virus, endogenous feline retrovirus (*e.g.*, RD114), and mouse or rat gL30 sequences used as a retroviral vector. Particularly preferred strains of MLV from which recombinant retroviruses can be generated include 4070A and 1504A (Hartley and Rowe, *J. Vir.* 19:19, 1976), Abelson (ATCC No. VR-999), Friend (ATCC No. VR-245), Graffi (Ru *et al.*, *J. Vir.* 67:4722, 1993; and Yantchev *Neoplasma* 26:397, 1979), Gross (ATCC No.

VR-590), Kirsten (Albino *et al.*, *J. Exp. Med.* 164:1710, 1986), Harvey sarcoma virus (Manly *et al.*, *J. Vir.* 62:3540, 1988; and Albino *et al.*, *J. Exp. Med.* 164:1710, 1986) and Rauscher (ATCC No. VR-998), and Moloney MLV (ATCC No. VR-190). A particularly preferred non-mouse retrovirus is Rous sarcoma virus. Preferred Rous sarcoma viruses include Bratislava (Manly *et al.*, *J. Vir.* 62:3540, 1988; and Albino *et al.*, *J. Exp. Med.* 164:1710, 1986), Bryan high titer (*e.g.*, ATCC Nos. VR-334, VR-657, VR-726, VR-659, and VR-728), Bryan standard (ATCC No. VR-140), Carr-Zilber (Adighitov *et al.*, *Neoplasma* 27:159, 1980), Engelbreth-Holm (Laurent *et al.*, *Biochem Biophys Acta* 908:241, 1987), Harris, Prague (*e.g.*, ATCC Nos. VR-772, and 45033), and Schmidt-Ruppin (*e.g.* ATCC Nos. VR-724, VR-725, VR-354) viruses.

Any of the above retroviruses can be readily utilized in order to assemble or construct retroviral gene delivery vehicles given the disclosure provided herein and standard recombinant techniques (*e.g.*, Sambrook *et al.*, 1989, and Kunkle, *Proc. Natl. Acad. Sci. U.S.A.* 82:488, 1985) known in the art. Portions of retroviral expression vectors can be derived from different retroviruses. For example, retrovector LTRs can be derived from a murine sarcoma virus, a tRNA binding site from a Rous sarcoma virus, a packaging signal from a murine leukemia virus, and an origin of second strand synthesis from an avian leukosis virus. These recombinant retroviral vectors can be used to generate transduction competent retroviral vector particles by introducing them into appropriate packaging cell lines (*see* Serial No. 07/800,921, filed November 29, 1991). Recombinant retroviruses can be produced which direct the site-specific integration of the recombinant retroviral genome into specific regions of the host cell DNA. Such site-specific integration can be mediated by a chimeric integrase incorporated into the retroviral particle (*see* Serial No. 08/445,466 filed May 22, 1995). It is preferable that the recombinant viral gene delivery vehicle is a replication-defective recombinant virus.

Packaging cell lines suitable for use with the above-described retroviral gene delivery vehicles can be readily prepared (*see* Serial No. 08/240,030, filed May 9, 1994; *see also* WO 92/05266) and used to create producer cell lines (also termed vector cell lines or "VCLs") for production of recombinant viral particles. In particularly preferred embodiments of the present invention, packaging cell lines are made from human (*e.g.*,

HT1080 cells) or mink parent cell lines, thereby allowing production of recombinant retroviral gene delivery vehicles which are capable of surviving inactivation in human serum. The construction of recombinant retroviral gene delivery vehicles is described in detail in WO 91/02805. These recombinant retroviral gene delivery vehicles can be used to generate transduction competent retroviral particles by introducing them into appropriate packaging cell lines (*see* Serial No. 07/800,921). Similarly, adenovirus gene delivery vehicles can also be readily prepared and utilized given the disclosure provided herein (*see also* Berkner, *Biotechniques* 6:616-627, 1988, and Rosenfeld *et al.*, *Science* 252:431-434, 1991, WO 93/07283, WO 93/06223, and WO 93/07282).

A gene delivery vehicle can also be a recombinant adenoviral gene delivery vehicle. Such vehicles can be readily prepared and utilized given the disclosure provided herein (*see* Berkner, *Biotechniques* 6:616, 1988, and Rosenfeld *et al.*, *Science* 252:431, 1991, WO 93/07283, WO 93/06223, and WO 93/07282). Adeno-associated viral gene delivery vehicles can also be constructed and used to deliver proteins or polynucleotides of the invention to cells *in vitro* or *in vivo*. The use of adeno-associated viral gene delivery vehicles *in vitro* is described in Chatterjee *et al.*, *Science* 258: 1485-1488 (1992), Walsh *et al.*, *Proc. Nat'l. Acad. Sci.* 89: 7257-7261 (1992), Walsh *et al.*, *J. Clin. Invest.* 94: 1440-1448 (1994), Flotte *et al.*, *J. Biol. Chem.* 268: 3781-3790 (1993), Ponnazhagan *et al.*, *J. Exp. Med.* 179: 733-738 (1994), Miller *et al.*, *Proc. Nat'l Acad. Sci.* 91: 10183-10187 (1994), Einerhand *et al.*, *Gene Ther.* 2: 336-343 (1995), Luo *et al.*, *Exp. Hematol.* 23: 1261-1267 (1995), and Zhou *et al.*, *Gene Therapy* 3: 223-229 (1996). *In vivo* use of these vehicles is described in Flotte *et al.*, *Proc. Nat'l Acad. Sci.* 90: 10613-10617 (1993), and Kaplitt *et al.*, *Nature Genet.* 8:148-153 (1994).

In another embodiment of the invention, a gene delivery vehicle is derived from a togavirus. Preferred togaviruses include alphaviruses, in particular those described in U.S. Serial No. 08/405,627, filed March 15, 1995, WO 95/07994. Alpha viruses, including Sindbis and ELVS viruses can be gene delivery vehicles for polynucleotides of the invention. Alpha viruses are described in WO 94/21792, WO 92/10578 and WO 95/07994. Several different alphavirus gene delivery vehicle systems can be constructed and used to deliver polynucleotides to a cell according to the present invention.

Representative examples of such systems include those described in U.S. Patents 5,091,309 and 5,217,879. Particularly preferred alphavirus gene delivery vehicles for use in the present invention include those which are described in WO 95/07994, and U.S. Serial No. 08/405,627.

5            Preferably, the recombinant viral vehicle is a recombinant alphavirus viral vehicle based on a Sindbis virus. Sindbis constructs, as well as numerous similar constructs, can be readily prepared essentially as described in U.S. Serial No. 08/198,450. Sindbis viral gene delivery vehicles typically comprise a 5' sequence capable of initiating Sindbis virus transcription, a nucleotide sequence encoding Sindbis non-structural proteins, a  
10           viral junction region inactivated so as to prevent fragment transcription, and a Sindbis RNA polymerase recognition sequence. Optionally, the viral junction region can be modified so that polynucleotide transcription is reduced, increased, or maintained. As will be appreciated by those in the art, corresponding regions from other alphaviruses can be used in place of those described above.

15           The viral junction region of an alphavirus-derived gene delivery vehicle can comprise a first viral junction region which has been inactivated in order to prevent transcription of the polynucleotide and a second viral junction region which has been modified such that polynucleotide transcription is reduced. An alphavirus-derived vehicle can also include a 5' promoter capable of initiating synthesis of viral RNA from  
20           cDNA and a 3' sequence which controls transcription termination.

             Other recombinant togaviral gene delivery vehicles which can be utilized in the present invention include those derived from Semliki Forest virus (ATCC VR-67; ATCC VR-1247), Middleberg virus (ATCC VR-370), Ross River virus (ATCC VR-373; ATCC VR-1246), Venezuelan equine encephalitis virus (ATCC VR923; ATCC VR-1250;  
25           ATCC VR-1249; ATCC VR-532), and those described in U.S. Patents 5,091,309 and 5,217,879 and in WO 92/10578. The Sindbis vehicles described above, as well as numerous similar constructs, can be readily prepared essentially as described in U.S. Serial No. 08/198,450.

             Other viral gene delivery vehicles suitable for use in the present invention  
30           include, for example, those derived from poliovirus (Evans *et al.*, *Nature* 339:385, 1989,

and Sabin *et al.*, *J. Biol. Standardization* 1:115, 1973) (ATCC VR-58); rhinovirus (Arnold *et al.*, *J. Cell. Biochem.* L401, 1990) (ATCC VR-1110); pox viruses, such as canary pox virus or vaccinia virus (Fisher-Hoch *et al.*, *PROC. NATL. ACAD. SCI. U.S.A.* 86:317, 1989; Flexner *et al.*, *Ann. N.Y. Acad. Sci.* 569:86, 1989; Flexner *et al.*, *Vaccine* 8:17, 1990; U.S. 4,603,112 and U.S. 4,769,330; WO 89/01973) (ATCC VR-111; ATCC VR-2010); SV40 (Mulligan *et al.*, *Nature* 277:108, 1979) (ATCC VR-305), (Madzak *et al.*, *J. Gen. Vir.* 73:1533, 1992); influenza virus (Luytjes *et al.*, *Cell* 59:1107, 1989; McMichael *et al.*, *The New England Journal of Medicine* 309:13, 1983; and Yap *et al.*, *Nature* 273:238, 1978) (ATCC VR-797); parvovirus such as adeno-associated virus (Samulski *et al.*, *J. Vir.* 63:3822, 1989, and Mendelson *et al.*, *Virology* 166:154, 1988) (ATCC VR-645); herpes simplex virus (Kit *et al.*, *Adv. Exp. Med. Biol.* 215:219, 1989) (ATCC VR-977; ATCC VR-260); *Nature* 277: 108, 1979); human immunodeficiency virus (EPO 386,882, Buchschacher *et al.*, *J. Vir.* 66:2731, 1992); measles virus (EPO 440,219) (ATCC VR-24); A (ATCC VR-67; ATCC VR-1247), Aura (ATCC VR-368), Bebaru virus (ATCC VR-600; ATCC VR-1240), Cabassou (ATCC VR-922), Chikungunya virus (ATCC VR-64; ATCC VR-1241), Fort Morgan (ATCC VR-924), Getah virus (ATCC VR-369; ATCC VR-1243), Kyzylagach (ATCC VR-927), Mayaro (ATCC VR-66), Mucambo virus (ATCC VR-580; ATCC VR-1244), Ndumu (ATCC VR-371), Pixuna virus (ATCC VR-372; ATCC VR-1245), Tonate (ATCC VR-925), Trinititi (ATCC VR-469), Una (ATCC VR-374), Whataroa (ATCC VR-926), Y-62-33 (ATCC VR-375), O'Nyong virus, Eastern encephalitis virus (ATCC VR-65; ATCC VR-1242), Western encephalitis virus (ATCC VR-70; ATCC VR-1251; ATCC VR-622; ATCC VR-1252), and coronavirus (Hamre *et al.*, *Proc. Soc. Exp. Biol. Med.* 121:190, 1966) (ATCC VR-740).

25           A polynucleotide of the invention can also be combined with a condensing agent to form a gene delivery vehicle. In a preferred embodiment, the condensing agent is a polycation, such as polylysine, polyarginine, polyornithine, protamine, spermine, spermidine, and putrescine. Many suitable methods for making such linkages are known in the art (see, for example, Serial No. 08/366,787, filed December 30, 1994).

30           In an alternative embodiment, a polynucleotide is associated with a liposome to

form a gene delivery vehicle. Liposomes are small, lipid vesicles comprised of an aqueous compartment enclosed by a lipid bilayer, typically spherical or slightly elongated structures several hundred Angstroms in diameter. Under appropriate conditions, a liposome can fuse with the plasma membrane of a cell or with the  
5 membrane of an endocytic vesicle within a cell which has internalized the liposome, thereby releasing its contents into the cytoplasm. Prior to interaction with the surface of a cell, however, the liposome membrane acts as a relatively impermeable barrier which sequesters and protects its contents, for example, from degradative enzymes. Additionally, because a liposome is a synthetic structure, specially designed liposomes  
10 can be produced which incorporate desirable features. See Stryer, *Biochemistry*, pp. 236-240, 1975 (W.H. Freeman, San Francisco, CA); Szoka *et al.*, *Biochim. Biophys. Acta* 600:1, 1980; Bayer *et al.*, *Biochim. Biophys. Acta* 550:464, 1979; Rivnay *et al.*, *Meth. Enzymol.* 149:119, 1987; Wang *et al.*, *PROC. NATL. ACAD. SCI. U.S.A.* 84: 7851, 1987, Plant *et al.*, *Anal. Biochem.* 176:420, 1989, and U.S. Patent 4,762,915. Liposomes can  
15 encapsulate a variety of nucleic acid molecules including DNA, RNA, plasmids, and expression constructs comprising polynucleotides such those disclosed in the present invention.

Liposomal preparations for use in the present invention include cationic (positively charged), anionic (negatively charged) and neutral preparations. Cationic  
20 liposomes have been shown to mediate intracellular delivery of plasmid DNA (Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7413-7416, 1987), mRNA (Malone *et al.*, *Proc. Natl. Acad. Sci. USA* 86:6077-6081, 1989), and purified transcription factors (Debs *et al.*, *J. Biol. Chem.* 265:10189-10192, 1990), in functional form. Cationic liposomes are readily available. For example, N[1-2,3-dioleoyloxy)propyl]-N,N,N-triethylammonium  
25 (DOTMA) liposomes are available under the trademark Lipofectin, from GIBCO BRL, Grand Island, NY. See also Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 91: 5148-5152.87, 1994. Other commercially available liposomes include Transfectace (DDAB/DOPE) and DOTAP/DOPE (Boehringer). Other cationic liposomes can be prepared from readily available materials using techniques well known in the art. See, e.g., Szoka *et al.*, *Proc.*  
30 *Natl. Acad. Sci. USA* 75:4194-4198, 1978; and WO 90/11092 for descriptions of the

synthesis of DOTAP (1,2-bis(oleoyloxy)-3-(trimethylammonio)propane) liposomes.

Similarly, anionic and neutral liposomes are readily available, such as from Avanti Polar Lipids (Birmingham, AL), or can be easily prepared using readily available materials. Such materials include phosphatidyl choline, cholesterol, phosphatidyl  
5 ethanolamine, dioleoylphosphatidyl choline (DOPC), dioleoylphosphatidyl glycerol (DOPG), dioleoylphosphatidyl ethanolamine (DOPE), among others. These materials can also be mixed with the DOTMA and DOTAP starting materials in appropriate ratios. Methods for making liposomes using these materials are well known in the art.

The liposomes can comprise multilammellar vesicles (MLVs), small unilamellar  
10 vesicles (SUVs), or large unilamellar vesicles (LUVs). The various liposome-nucleic acid complexes are prepared using methods known in the art. *See, e.g.,* Straubinger *et al.*, *METHODS OF IMMUNOLOGY* (1983), Vol. 101, pp. 512-527; Szoka *et al.*, *Proc. Natl. Acad. Sci. USA* 87:3410-3414, 1990; Papahadjopoulos *et al.*, *Biochim. Biophys. Acta* 394:483, 1975; Wilson *et al.*, *Cell* 17:77, 1979; Deamer and Bangham, *Biochim.*  
15 *Biophys. Acta* 443:629, 1976; Ostro *et al.*, *Biochem. Biophys. Res. Commun.* 76:836, 1977; Fraley *et al.*, *Proc. Natl. Acad. Sci. USA* 76:3348, 1979; Enoch and Strittmatter, *Proc. Natl. Acad. Sci. USA* 76:145, 1979; Fraley *et al.*, *J. Biol. Chem.* 255:10431, 1980; Szoka and Papahadjopoulos, *Proc. Natl. Acad. Sci. USA* 75:145, 1979; and Schaefer-Ridder *et al.*, *Science* 215:166, 1982.

20 In addition, lipoproteins can be included with a polynucleotide of the invention for delivery to a cell. Examples of such lipoproteins include chylomicrons, HDL, IDL, LDL, and VLDL. Mutants, fragments, or fusions of these proteins can also be used. Modifications of naturally occurring lipoproteins can also be used, such as acetylated LDL. These lipoproteins can target the delivery of polynucleotides to cells expressing  
25 lipoprotein receptors. Preferably, if lipoproteins are included with a polynucleotide, no other targeting ligand is included in the composition.

In another embodiment, naked polynucleotide molecules are used as gene delivery vehicles, as described in WO 90/11092 and U.S. Patent 5,580,859. Such gene delivery vehicles can be either DNA or RNA and, in certain embodiments, are linked to  
30 killed adenovirus. Curiel *et al.*, *Hum. Gene. Ther.* 3:147-154, 1992. Other suitable

vehicles include DNA-ligand (Wu *et al.*, *J. Biol. Chem.* 264:16985-16987, 1989), lipid-DNA combinations (Felgner *et al.*, *Proc. Natl. Acad. Sci. USA* 84:7413 7417, 1989), liposomes (Wang *et al.*, *Proc. Natl. Acad. Sci.* 84:7851-7855, 1987) and microprojectiles (Williams *et al.*, *Proc. Natl. Acad. Sci.* 88:2726-2730, 1991).

5           One can increase the efficiency of naked polynucleotide uptake into cells by coating the polynucleotides onto biodegradable latex beads. This approach takes advantage of the observation that latex beads, when incubated with cells in culture, are efficiently transported and concentrated in the perinuclear region of the cells. The beads will then be transported into cells when injected into muscle. Polynucleotide-coated  
10 latex beads will be efficiently transported into cells after endocytosis is initiated by the latex beads and thus increase gene transfer and expression efficiency. This method can be improved further by treating the beads to increase their hydrophobicity, thereby facilitating the disruption of the endosome and release of polynucleotides into the cytoplasm.

15           One polynucleotide of the invention is designated hCornichon. The nucleotide sequence of hCornichon is shown in SEQ ID NO:1. hCornichon cDNA represents a transcript of 1325 nucleotides with a translation stop codon (TAG) at position 428, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at position 1292, and a poly(A) tail at position 1316. The DNA sequence between nucleotides 2 and 427 encodes a protein  
20 of 142 amino acids, as shown in SEQ ID NO:2. A potential signal peptide is located in the first 28 amino acid residues. An hCornichon polynucleotide can comprise at least 499, 550, 600, 700, 750, 800, 850, 850, 900, 950, 1000, 1100, 1141, 1150, 1200, or 1250 nucleotides of SEQ ID NO:1 or the complements thereof.

          Another polynucleotide of the invention is designated BMS46. The nucleotide  
25 sequence of BMS46 is shown in SEQ ID NO:3. BMS46 cDNA represents a transcript of 1277 nucleotides with a translation start codon (ATG) at position 656, a translation stop codon (TAG) at position 1223, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at position 1243, and a poly(A) tail at position 1260. The DNA sequence between nucleotides 656 and 1222 encodes a protein of 189 amino acid residues, as shown in  
30 SEQ ID NO:4. A potential signal peptide is located in the first 47 amino acid residues.



A BMS46 polynucleotide can comprise at least 474, 475, 476, 477, 500, 550, 600, 650, 700, 750, 800, 850, 900, 950, 1000, 1050, 1150, 1200, or 1250 contiguous nucleotides of SEQ ID NO:3, or at least 313, 314, 315, or 316 contiguous nucleotides selected from nucleotides 1-1001 of SEQ ID NO:3, or the complements thereof.

5           The nucleotide sequence of another polynucleotide of the invention, termed BMS112, is shown in SEQ ID NO:5. BMS112 cDNA represents a transcript of 1610 nucleotides with a translation start codon (ATG) at position 132, a translation stop codon (TGA) at position 1251, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at position 1516, and a poly(A) tail at position 1594. The DNA sequence between  
10   nucleotides 132 and 1250 encodes a polypeptide of 373 amino acid residues (SEQ ID NO:6). A BMS112 polynucleotide can comprise at least 538, 600, 700, 751, 800, 850, 900, 950, 1000, 1200, 1300, 1400, 1500 or 1600 contiguous nucleotides of SEQ ID NO:5, at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides selected from nucleotides 1-946, at least 13 contiguous nucleotides selected from nucleotides 1-  
15   1039 of SEQ ID NO:5, or the complements thereof.

Yet another polynucleotide of the invention has the nucleotide sequence shown in SEQ ID NO:7 and is designated BMS118. BMS118 cDNA represents a transcript of 1499 nucleotides with a translation start codon (ATG) at position 140, a translation stop codon (TAA) at position 1358, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at  
20   position 1463, and a poly(A) tail at position 1482. The DNA sequence between nucleotides 140 and 1357 encodes a polypeptide of 406 amino acid residues (SEQ ID NO:8). The potential signal peptide of the BMS118 protein is located in the first 29 amino acids. A BMS118 polynucleotide can comprise at least 522, 550, 600, 651, 700, 750, 800, 850, 900, 950, 1000, 1050, 1100, 1150, 1200, 1250, 1300, 1350, 1400, or 1450  
25   contiguous nucleotides of SEQ ID NO:7, at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides selected from nucleotides 1-913 of SEQ ID NO:7, or the complements thereof.

Another polynucleotide of the invention has the nucleotide sequence shown in SEQ ID NO:9 and is designated BMS164. BMS164 cDNA represents a transcript of  
30   1272 nucleotides with a translation start codon (ATG) at position 313 and a translation

stop codon (TAG) at position 1186. The DNA sequence between nucleotides 313 and 1185 encodes a polypeptide of 291 amino acid residues (SEQ ID NO:10). A BMS164 polynucleotide can comprise at least 317, 400, 484, 500, 600, 700, 800, 900, 1000, 1100, or 1200 contiguous nucleotides of SEQ ID NO:9, at least 183 contiguous nucleotides  
5 selected from nucleotides 1-984 of SEQ ID NO:9, or at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides selected from nucleotides 1-216 or 379-812 of SEQ ID NO:9, or the complements thereof.

Another polynucleotide of the invention, BMS192, has the nucleotide sequence shown in SEQ ID NO:11. BMS192 cDNA represents a transcript of 1585 nucleotides  
10 with a translation start codon (ATG) at position 41, a translation stop codon (TGA) at position 1190, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at position 1439, and a poly(A) tail at position 1574. The DNA sequence between nucleotides 41 and 1189 encodes a polypeptide of 383 amino acid residues (SEQ ID NO:12). The potential signal peptide of the BMS192 protein is located in the first 19 amino acids. A BMS192  
15 polynucleotide can comprise at least 289, 300, 400, 500, 594, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, or 1500 contiguous nucleotides of SEQ ID NO:11, at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides selected from nucleotides 1-585 or 853-1120 of SEQ ID NO:11, or the complements thereof.

Another polynucleotide of the invention, BMS227, has the nucleotide sequence shown in SEQ ID NO:13. BMS227 cDNA represents a transcript of 1071 nucleotides  
20 with a translation start codon (ATG) at position 151, a translation stop codon (TGA) at position 934, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at position 1018, and a poly(A) tail at position 1053. The DNA sequence between nucleotides 151 and 933 encodes a polypeptide of 261 amino acid residues (SEQ ID NO:14). The potential  
25 signal peptide of the BMS227 protein is located in the first 32 amino acids. A BMS227 polynucleotide can comprise 275, 300, 400, 500, 592, 600, 700, 800, 900, or 1000 contiguous nucleotides of SEQ ID NO: 13, at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides selected from nucleotides 1-294 of SEQ ID NO:13, or the complements thereof.

30 Yet another polynucleotide of the invention is designated BMS115. The

nucleotide sequence of BMS115 is shown in SEQ ID NO:15. BMS115 cDNA represents a transcript of 2520 nucleotides with a translation start codon (ATG) at position 1, a translation stop codon at position 1666, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at position 2470, and a poly(A) tail at position 2503. The DNA sequence  
5 between nucleotides 1 and 1665 encodes a protein of 555 amino acids, as shown in SEQ ID NO:16. A potential signal peptide is located in the first 31 amino acid residues. A BMS115 polynucleotide can comprise at least 537, 600, 700, 800, 900, 1000, 1250, 1500, 1750, 2000, 2250, or 2500 contiguous nucleotides of SEQ ID NO:15, at least 294 contiguous nucleotides selected from nucleotides 1-1889 of SEQ ID NO:15, at least 171  
10 contiguous nucleotides selected from nucleotides 318-1766 of SEQ ID NO:15, or at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides selected from nucleotides 1-42, 478-908, or 1059-1078 of SEQ ID NO:15, or the complements thereof.

Yet another polynucleotide of the invention is designated BMS143. The nucleotide sequence of BMS143 is shown in SEQ ID NO:17. BMS143 cDNA represents  
15 a transcript of 1245 nucleotides with a translation start codon (ATG) at position 89, a translation stop codon at position 785, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at position 1199, and a poly(A) tail at position 1231. The DNA sequence between nucleotides 89 and 784 encodes a protein of 232 amino acids, as shown in SEQ ID NO:18. A potential signal peptide is located in the first 54 amino acid residues. A  
20 BMS143 polynucleotide can comprise at least 205, 300, 400, 500, 600, 700, 800, 900, 1000, 1100, or 1200 contiguous nucleotides of SEQ ID NO:17, or the complements thereof.

Yet another polynucleotide of the invention is designated BMS155. The nucleotide sequence of BMS155 is shown in SEQ ID NO:19. BMS155 cDNA represents  
25 a transcript of 1030 nucleotides with a translation start codon (ATG) at position 4, a translation stop codon at position 451, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at position 987, and a poly(A) tail at position 1016. The DNA sequence between nucleotides 4 and 450 encodes a protein of 149 amino acids, as shown in SEQ ID NO:20. A potential signal peptide is located in the first 47 amino acid residues. A BMS155  
30 polynucleotide can comprise at least 440, 500, 600, 700, 800, 900, or 1000 contiguous

nucleotides of SEQ ID NO:19 or the complements thereof.

Yet another polynucleotide of the invention is designated BMS208. The nucleotide sequence of BMS208 is shown in SEQ ID NO:21. BMS208 cDNA represents a transcript of 1563 nucleotides with a translation start codon (ATG) at position 255, a translation stop codon at position 756, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at position 1531, and a poly(A) tail at position 1550. The DNA sequence between nucleotides 255 and 755 encodes a protein of 167 amino acids, as shown in SEQ ID NO:22. A potential signal peptide is located in the first 62 amino acid residues. A BMS208 polynucleotide can comprise at least 451, 500, 600, 750, 1000, 1250, or 1500 contiguous nucleotides of SEQ ID NO:21, at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides selected from nucleotides 1-121 or 474-592 of SEQ ID NO:21, or the complements thereof.

Yet another polynucleotide of the invention is designated BMS235. The nucleotide sequence of BMS235 is shown in SEQ ID NO:23. BMS235 cDNA represents a transcript of 2590 nucleotides with a translation start codon (ATG) at position 29, a translation stop codon at position 872, and a poly(A) tail at position 1526. The DNA sequence between nucleotides 29 and 871 encodes a protein of 281 amino acids, as shown in SEQ ID NO:24. A potential signal peptide is located in the first 25 amino acid residues. A BMS235 polynucleotide can comprise at least 351 contiguous nucleotides of SEQ ID NO:23, at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides selected from nucleotides 1-612, 611-719, 713-830, or 830-1933 of SEQ ID NO:23, at least 21 contiguous nucleotides selected from nucleotides 1-1943 of SEQ ID NO:23, or the complements thereof.

Yet another polynucleotide of the invention is designated BMS240. The nucleotide sequence of BMS240 is shown in SEQ ID NO:25. BMS240 cDNA represents a transcript of 1668 nucleotides with a translation start codon (ATG) at position 99, a translation stop codon at position 807, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at position 1626, and a poly(A) tail at position 1655. The DNA sequence between nucleotides 99 and 806 encodes a protein of 236 amino acids, as shown in SEQ ID NO:26. A BMS240 polynucleotide can comprise at least 492, 500, 600, 750, 1000,

1250, 1500, or 1600 contiguous nucleotides of SEQ ID NO:25, at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides selected from nucleotides 758-847 of SEQ ID NO:25, or the complements thereof.

Yet another polynucleotide of the invention is designated BMS53. The  
5 nucleotide sequence of BMS53 is shown in SEQ ID NO:27. BMS53 cDNA represents a transcript of 1697 nucleotides with a translation start codon (ATG) at position 29, a translation stop codon at position 1427, a polyadenylation signal (ATTAAA) (SEQ ID NO:46) at position 1659, and a poly(A) tail at position 1682. The DNA sequence between nucleotides 29 and 1426 encodes a polypeptide of 466 amino acid residues, as  
10 shown in SEQ ID NO:28. A BMS53 polynucleotide can comprise at least 1024, 1100, 1200, 1300, 1400, 1500, or 1600 contiguous nucleotide of SEQ ID NO:27 or the complements thereof.

Yet another polynucleotide of the invention is designated BMS100. The nucleotide sequence of BMS100 is shown in SEQ ID NO:29. BMS100 cDNA represents  
15 a transcript of 1830 nucleotides with a translation start codon (ATG) at position 218, a translation stop codon at position 851, a polyadenylation signal (AATAAA) (SEQ ID NO:35) at position 1792, and a poly(A) tail at position 1811. The DNA sequence between nucleotides 218 and 850 encodes a protein of 211 amino acids, as shown in SEQ ID NO:30. A potential signal peptide is located in the first 18 amino acid residues. A  
20 BMS100 polynucleotide can comprise at least 347, 400, 500, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, 1500, 1600, 1700, or 1800 contiguous nucleotides of SEQ ID NO:29, at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides selected from nucleotides 548-601 of SEQ ID NO:29, or the complements thereof.

Yet another polynucleotide of the invention is designated BMS199. The  
25 nucleotide sequence of BMS199 is shown in SEQ ID NO:31. BMS199 cDNA represents a transcript of 1102 nucleotides with a translation start codon (ATG) at position 267, a translation stop codon at position 990, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at position 1072, and a poly(A) tail at position 1089. The DNA sequence between nucleotides 267 and 989 encodes a protein of 241 amino acids, as shown in SEQ  
30 ID NO:32. A potential signal peptide is located in the first 32 amino acid residues. A

BMS199 polynucleotide can comprise at least 394, 400, 500, 600, 700, 800, 900, 1000, or 1100 contiguous nucleotides of SEQ ID NO:31, at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides selected from nucleotides 1-361 or 1083-1102 of SEQ ID NO:31, or the complements thereof.

5           Yet another polynucleotide of the invention is designated BMS206. The nucleotide sequence of BMS206 is shown in SEQ ID NO:33. BMS206 cDNA represents a transcript of 966 nucleotides with a translation start codon (ATG) at position 36, a translation stop codon at position 585, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at position 920, and a poly(A) tail at position 949. The DNA sequence between  
10       nucleotides 36 and 584 encodes a protein of 183 amino acids, as shown in SEQ ID NO:34. A BMS206 polynucleotide can comprise at least 492, 500, 600, 700, 800, or 900 contiguous nucleotides of SEQ ID NO:33 or the complements thereof.

          Yet another polynucleotide of the invention is designated BMS242. The nucleotide sequence of BMS242 is shown in SEQ ID NO:35. BMS242 cDNA represents  
15       a transcript of 1570 nucleotides with a translation start codon (ATG) at position 76, a translation stop codon at position 1030, and a poly (1) tail at position 1562. The DNA sequence between nucleotides 76 and 1029 encodes a protein of 318 amino acid residues, as shown in SEQ ID NO:36. A BMS242 polynucleotide can comprise at least 510, 600, 700, 800, 900, 1000, 1100, 1200, 1300, 1400, or 1500 contiguous nucleotides of SEQ ID  
20       NO:35, at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides selected from nucleotides 1-502 or 505-631 of SEQ ID NO:35, or the complements thereof.

          Yet another polynucleotide of the invention is termed BMS37. The nucleotide sequence of BMS37 is shown in SEQ ID NO:37. BMS37 cDNA represents a transcript  
25       of 1542 nucleotides with a translation start codon (ATG) at position 121, a translation stop codon at position 1105, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at position 1508, and a poly(A) tail at position 1526. The DNA sequence between nucleotides 121 and 1104 encodes a protein of 328 amino acid residues, as shown in SEQ ID NO:38. The potential signal peptide the BMS37 protein is located in the first 20  
30       amino acids. A BMS37 polynucleotide can comprise at least 392, 400, 500, 600, 700,

800, 900, 1000, 1100, 1200, 1300, 1400, or 1500 contiguous nucleotides of SEQ ID NO:37, at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides selected from nucleotides 1-502 or 505-631 of SEQ ID NO:37, or the complements thereof.

5 Yet another polynucleotide of the invention is designated BMS42. The nucleotide sequence of BMS42 is shown in SEQ ID NO:39. BMS42 cDNA represents a transcript of 1990 nucleotides with a translation start codon (ATG) at position 104, a translation stop codon at position 1615, a polyadenylation signal (AATAAA) (SEQ ID NO:45) at position 1952, and a poly(A) tail at position 1971. The DNA sequence  
10 between nucleotides 104 and 1614 encodes a protein of 504 amino acids, as shown in SEQ ID NO:40. A potential signal peptide is located in the first 67 amino acids. A BMS42 polynucleotides can comprise at least 559, 600, 700, 800, 900, 10000, 1250, 1500, 1750, 1800, or 1900 contiguous nucleotides of SEQ ID NO:39, at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides selected from nucleotides 1-92 of  
15 SEQ ID NO:39, or the complements thereof.

Yet another polynucleotide of the invention is designated BMS60. The nucleotide sequence of BMS60 is shown in SEQ ID NO:41. BMS60 cDNA represents a transcript of 684 nucleotides with a translation start codon (ATG) at position 7, a translation stop codon at position 445, a polyadenylation signal (AATAAA) (SEQ ID  
20 NO:45) at position 644, and a poly(A) tail at position 667. The DNA sequence between nucleotides 7 and 444 encodes a protein of 146 amino acid residues, as shown in SEQ ID NO:42. A potential signal peptide is located in the first 20 amino acids. A BMS60 polynucleotide can comprise at least 254, 300, 350, 400, 450, 500, 550, 600, or 650 contiguous nucleotides of SEQ ID NO:41, at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40,  
25 or 50 contiguous nucleotides selected from nucleotides 1-34 or 55-110 of SEQ ID NO:41, or the complements thereof.

Yet another polynucleotide of the invention is designated BMS61. The nucleotide sequence of BMS61 is shown in SEQ ID NO:43. BMS61 cDNA represents a transcript of 1152 nucleotide with a translation start codon (ATG) at position 276, a  
30 translation stop codon at position 795, and a poly(A) tail at position 1150. The DNA

sequence between nucleotides 276 and 794 encodes a protein of 173 amino acid residues, as shown in SEQ ID NO:44. A BMS61 polynucleotide can comprise at least 103, 200, 300, 400, 500, 600, 700, 800, 900, 1000 or 1100 contiguous nucleotides of SEQ ID NO:43, at least 11, 12, 13, 14, 15, 20, 25, 30, 35, 40, or 50 contiguous nucleotides  
5 selected from nucleotides 1-280, 270-319, 378-423, 414-492, 532-570, or 1086-1152 of SEQ ID NO:43, or the complements thereof.

The present invention provides isolated genes which comprise the coding sequences disclosed herein. The genes can be isolated in accordance with known methods using the sequence information disclosed herein. Such methods include the  
10 preparation of probes or primers from the disclosed sequence information for identification and/or amplification of genes in appropriate genomic libraries or other sources of genomic materials.

The invention also provides means of altering the expression of genes which have the coding sequences disclosed herein. In one embodiment of the invention, expression  
15 of an endogenous gene having a coding sequence as shown in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, or 43 in a cell can be altered by introducing in frame with the endogenous gene a DNA construct comprising a transcription unit by homologous recombination to form a homologously recombinant cell comprising the transcription unit. The transcription unit comprises a targeting  
20 sequence, a regulatory sequence, an exon, and an unpaired splice donor site. This method of affecting endogenous gene expression is taught in U.S. Patent No. 5,641,670.

The targeting sequence is a segment of at least 10, 12, 15, 20, or 50 contiguous nucleotides selected from the nucleotide sequences shown in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, and 43. The transcription  
25 unit is located upstream to a coding sequence of the endogenous gene. The exogenous regulatory sequence directs transcription of the coding sequence of the gene.

In another embodiment of the invention, expression of a gene with a coding sequence as shown in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, or 43 is decreased using a ribozyme, an RNA molecule with  
30 catalytic activity. *See, e.g., Cech, 1987, Science 236: 1532-1539; Cech, 1990, Ann.*



*Rev. Biochem.* 59:543-568; Cech, 1992, *Curr. Opin. Struct. Biol.* 2: 605-609; Couture and Stinchcomb, 1996, *Trends Genet.* 12: 510-515. Ribozymes can be used to inhibit gene function by cleaving an RNA sequence, as is known in the art (*e.g.*, Haseloff *et al.*, U.S. 5,641,673).

5           The coding sequences disclosed herein can be used to generate a ribozyme which will specifically bind to the corresponding mRNA. Methods of designing and constructing ribozymes which can cleave other RNA molecules in trans in a highly sequence specific manner have been developed and described in the art (see Haseloff *et al.*, *Nature* 334:585-591, 1988). For example, the cleavage activity of ribozymes can be  
10           targeted to specific RNAs by engineering a discrete "hybridization" region into the ribozyme. The hybridization region contains a sequence complementary to the target RNA and thus specifically hybridizes with the target (see, for example, Gerlach *et al.*, EP 321,201). Longer complementary sequences can be used to increase the affinity of the hybridization sequence for the target. The hybridizing and cleavage regions of the  
15           ribozyme can be integrally related; thus, upon hybridizing to the target RNA through the complementary regions, the catalytic region of the ribozyme can cleave the target.

          Ribozymes can be introduced into cells as part of a DNA construct, as is known in the art. The DNA construct can also include transcriptional regulatory elements, such as a promoter element, an enhancer or UAS element, and a transcriptional terminator  
20           signal, for controlling transcription of the ribozyme in the cells.

          Mechanical methods, such as microinjection, liposome-mediated transfection, electroporation, or calcium phosphate precipitation, can be used to introduce the ribozyme-containing DNA construct into cells in order to decrease gene expression. Alternatively, if it is desired that the cells stably retain the DNA construct, it can be  
25           supplied on a plasmid and maintained as a separate element or integrated into the genome of the cells, as is known in the art.

          Expression of a gene with a coding sequence as shown in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, or 43 can also be altered using an antisense oligonucleotide. The sequence of the antisense oligonucleotide is  
30           complementary to at least a portion of a coding sequence disclosed herein. Preferably,

the antisense oligonucleotide is at least six nucleotides in length, but can be at least 8, 11, 12, 15, 20, 25, 30, 35, 40, 45, or 50 nucleotides long. Longer sequences, such as the complement of the nucleotide sequences shown in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, or 43, can also be used. Antisense  
5 oligonucleotides can be provided in a construct of the invention and introduced into cells using transfection techniques known in the art.

Antisense oligonucleotides can be composed of deoxyribonucleotides, ribonucleotides, or a combination of both. Oligonucleotides can be synthesized manually or by an automated synthesizer, by covalently linking the 5' end of one nucleotide with  
10 the 3' end of another nucleotide with non-phosphodiester internucleotide linkages such as alkylphosphonates, phosphorothioates, phosphorodithioates, alkylphosphonothioates, alkylphosphonates, phosphoramidates, phosphate esters, carbamates, acetamidate, carboxymethyl esters, carbonates, and phosphate triesters. See Brown, 1994, *Meth. Mol. Biol.* 20:1-8; Sonveaux, 1994, *Meth. Mol. Biol.* 26:1-72; Uhlmann *et al.*, 1990, *Chem.*  
15 *Rev.* 90:543-583.

Precise complementarity is not required for successful duplex formation between an antisense molecule and its complementary coding sequence. Antisense molecules which comprise, for example, 2, 3, 4, or 5 or more stretches of contiguous nucleotides which are precisely complementary to a coding sequence of the invention, each separated  
20 by a stretch of contiguous nucleotides which are not complementary to adjacent coding sequences, can provide targeting specificity for mRNA. Preferably, each stretch of contiguous nucleotides is at least 4, 5, 6, 7, or 8 or more nucleotides in length. Non-complementary intervening sequences are preferably 1, 2, 3, or 4 nucleotides in length. One skilled in the art can easily use the calculated melting point of an antisense-sense  
25 pair to determine the degree of mismatching which will be tolerated between a particular antisense oligonucleotide and a particular coding sequence of the invention.

Antisense oligonucleotides can be modified without affecting their ability to hybridize to a coding sequence of the invention. These modifications can be internal or at one or both ends of the antisense oligonucleotide. For example, internucleoside  
30 phosphate linkages can be modified by adding cholesteryl or diamine moieties with

varying numbers of carbon residues between the amino groups and terminal ribose.

Modified bases and/or sugars, such as arabinose instead of ribose, or a 3', 5'-substituted oligonucleotide in which the 3' hydroxyl group or the 5' phosphate group are substituted, can also be employed in a modified antisense oligonucleotide. These modified

5 oligonucleotides can be prepared by methods well known in the art. Agrawal *et al.*, *Trends Biotechnol.* 10:152-158, 1992; Uhlmann *et al.*, *Chem. Rev.* 90:543-584, 1990; Uhlmann *et al.*, *Tetrahedron. Lett.* 215:3539-3542, 1987.

Antibodies of the invention can also be used to decrease the function of proteins of the invention. Specific antibodies bind to a protein of the invention to prevent the  
10 protein from functioning in the cell. Polynucleotides encoding single-chain antibodies of the invention can be introduced into cells using standard transfection techniques.

Alternatively, therapeutic antibodies of the invention can be targeted to a particular cell type, for example, by binding an antibody to a coupling molecule which is specific for both the antibody and the target, as disclosed in WO 95/08577. The coupling molecule  
15 can comprise immunoglobulin binding domains.

Proteins of the invention comprise the amino acid sequences shown in SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, and 44. Protein or polypeptide fragments which are capable of exhibiting biological activity are also encompassed by the present invention. Non-naturally  
20 occurring protein variants which retain substantially the same biological activities as naturally occurring proteins of the invention are also included here. Preferably, naturally or non-naturally occurring protein variants have amino acid sequences which are at least 65%, 75%, 85%, 90%, or 95% identical to the amino acid sequences shown in SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, and 44  
25 are secreted proteins, and have similar biological properties. More preferably, the molecules are 98% identical. Percent identity can be determined using computer programs which use the Smith-Waterman algorithm using an affine gap search with the following parameters: a gap open penalty of 12 and a gap extension penalty of 1.

Guidance in determining which amino acid residues may be substituted, inserted,  
30 or deleted without abolishing biological or immunological activity may be found using

computer programs well known in the art, such as DNASTAR software. Preferably, amino acid changes in protein variants or derivatives are conservative amino acid changes, *i.e.*, substitutions of similarly charged or uncharged amino acids. A conservative amino acid change involves substitution of one of a family of amino acids which are related in their side chains. Naturally occurring amino acids are generally divided into four families: acidic (aspartate, glutamate), basic (lysine, arginine, histidine), non-polar (alanine, valine, leucine, isoleucine, proline, phenylalanine, methionine, tryptophan), and uncharged polar (glycine, asparagine, glutamine, cystine, serine, threonine, tyrosine) amino acids. Phenylalanine, tryptophan, and tyrosine are sometimes classified jointly as aromatic amino acids. It is reasonable to expect that an isolated replacement of a leucine with an isoleucine or valine, an aspartate with a glutamate, a threonine with a serine, or a similar replacement of an amino acid with a structurally related amino acid will not have a major effect on the biological properties of the resulting protein variant.

15 Variants of proteins of the invention include glycosylated forms, aggregative conjugates with other molecules, and covalent conjugates with unrelated chemical moieties. Variants of the invention also include allelic variants, species variants, and muteins. Truncations or deletions of regions which do not affect the properties or functions of proteins of the invention are also variants. Covalent variants can be prepared by linkage of functionalities to groups which are found in the amino acid chain or at the N- or C-terminal residue, as is known in the art.

The invention also provides polypeptide fragments of the disclosed secreted proteins. Polypeptides of the invention comprise less than all of the amino acid sequences shown in SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, or 42 in the same primary order as found in the full-length amino acid sequences. For example, polypeptides of the invention can comprise at least 95, 100, 120, 130, or 140 contiguous amino acids of SEQ ID NO:2.

Other polypeptides of the invention can comprise at least 6, 8, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 101, 110, 120, 130, 150, 160, 170, or 180 contiguous amino acids of SEQ ID NO:4.

Yet other polypeptides of the invention can comprise at least 14, 15, 16, 18, 20, 25, or 30 contiguous amino acids selected from amino acids 1-312 of SEQ ID NO:6 or at least 75, 100, 125, 150, 175, 179, 200, 225, 250, 275, 300, 325, or 350 contiguous amino acids of SEQ ID NO:6.

5 Even other polypeptides of the invention can comprise at least 17, 18, 19, 20, 25, or 30 contiguous amino acids selected from amino acids 1-287 of SEQ ID NO:8 or at least 136, 140, 150, 150, 179, 200, 250, 300, 350, or 400 contiguous amino acids selected from SEQ ID NO:8.

Still other polypeptides of the invention can comprise at least 31, 32, 35, 40, or  
10 45 contiguous amino acids selected from amino acids 1-238 of SEQ ID NO:10 or at least 82, 85, 100, 132, 150, 200, 225, 250, or 275 contiguous amino acids of SEQ ID NO:10.

Other polypeptides of the invention can comprise at least 6, 7, 8, 9, 10, 15, or 20 contiguous amino acids selected from amino acids 1-184 or 270-362 of SEQ ID NO: 12, at least 8, 9, 10, 12, 15, 20, or 25 contiguous amino acids selected from amino acids 268-  
15 364 of SEQ ID NO: 12, at least 27, 30, 35, or 40 contiguous amino acids selected from amino acids 250-383 of SEQ ID NO:12, or at least 96, 100, 150, 200, 250, 300, or 350 contiguous amino acids selected from SEQ ID NO:12.

Yet other polypeptides of the invention can comprise at least 6, 7, 8, 9, 10, 12, 15, or 20 contiguous amino acids selected from amino acids 1-111 or 204-261 of SEQ ID  
20 NO: 14, at least 17, 18, 20, 25, or 30 contiguous amino acids selected from amino acids 1-150 of SEQ ID NO:14, at least 75, 80, 100, 104, 125, 150, 175, 200, 225, or 250 contiguous amino acids of SEQ ID NO:14.

Even other polypeptides of the invention can comprise at least 8, 10, 12, 14, 16, 20, 30, 40, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425,  
25 450, 475, 500, 525, or 550 contiguous amino acids of SEQ ID NO:16.

Still other polypeptides of the invention can comprise at least 39, 40, 45, 46, or 50 contiguous amino acids selected from amino acids 13-232 of SEQ ID NO:18 or at least 46, 50, 55, 60, 75, 100, 125, 150, 175, 200, or 225 contiguous amino acids of SEQ ID NO:18.

30 Other polypeptides of the invention can comprise at least 6, 8, 10, 12, 15, 20, 25,

30, 50, 75, 100, 125, or 140 contiguous amino acids from SEQ ID NO:20.

Yet other polypeptides of the invention can comprise at least 7, 8, 10, 12, 15, 20, 25, 30, 50, 75, 100, 125, 150, or 160 contiguous amino acids from SEQ ID NO:22.

Even other polypeptides of the invention comprise at least 7, 8, 10, 12, 15, 20, 25,  
5 30, 50, 75, 100, 125, 150, 175, 200, 225, 250, or 275 contiguous amino acids of SEQ ID NO:24.

Still other polypeptides of the invention comprise at least 11, 12, 15, 18, 20, 25, 30, 35, 50, 75, 100, 125, 150, 175, 200, or 225 contiguous amino acids of SEQ ID NO:26.

10 Other polypeptides of the invention comprise at least 6, 8, 10, 12, 15, or 20 contiguous amino acids selected from amino acids 1-31 of SEQ ID NO:28 or at least 257, 260, 270, 280, 290, 300, 325, 350, 375, 400, 425, or 450 contiguous amino acids of SEQ ID NO:28.

Yet other polypeptides of the invention comprise at least 6, 8, 10, 12, 15, 18, 20,  
15 25, 30, 50, 75, 100, 125, 150, 175, or 200 contiguous amino acids of SEQ ID NO:30.

Even other polypeptides of the invention comprise at least 6, 8, 10, 12, 15, or 20 contiguous amino acids selected from amino acids 1-65 of SEQ ID NO:32 or at least 117, 120, 150, 175, 200, or 225 contiguous amino acids of SEQ ID NO:32.

Still other polypeptides of the invention comprise at least 6, 8, 10, 12, 15, 20, 25,  
20 30, 50, 75, 100, 125, 150, or 175 contiguous amino acids of SEQ ID NO:34.

Other polypeptides of the invention comprise at least 14, 15, 18, 20, 25, 30, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, or 300 contiguous amino acids of SEQ ID NO:36.

Yet other polypeptides of the invention comprise at least 19, 20, 25, 30, 35, 40,  
25 50, 75, 100, 125, 150, 175, 200, 224, 250, 275, 300, or 325 contiguous amino acids of SEQ ID NO:38.

Even other polypeptides of the invention comprise at least 8, 10, 12, 15, 18, 20, 25, 30, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, or 500 contiguous amino acids of SEQ ID NO:40.

30 Still other polypeptides of the invention comprise at least 7, 8, 10, 12, 15, 20, 30,

50, 75, 100, or 125 contiguous amino acids of SEQ ID NO:42.

Other polypeptides of the invention comprise at least 10, 12, 15, 20, 25, 30, 50, 75, 100, 125, 150, or 170 contiguous amino acids of SEQ ID NO:44.

Polypeptides can be linear or can be cyclized using known methods, for example, as described in Saragovi *et al.*, *Bio/Technology* 10, 773-778 (1992) or McDowell *et al.*, *J. Amer. Chem. Soc.* 114, 9245-9253 (1992). Polypeptides can optionally be fused to carrier molecules such as immunoglobulins and used, for example, to increase the number of protein binding sites in a molecule or a molecular complex. Polypeptide fragments of the protein can be fused through linker sequences to the Fc portion of an immunoglobulin. Fusion of polypeptide fragments to the Fc portions of an IgG molecule can provide a bivalent form of a protein. Other immunoglobulin Fc portions, for example, IgM or IgA, can be used to provide multivalent forms of a protein.

Receptors or other membrane-bound proteins of the invention can be solubilized by deleting part of all of the intracellular and transmembrane domains of the protein, such that the protein can be fully secreted from a cell in which it is expressed. Intracellular and transmembrane domains of proteins of the invention can be identified using known techniques for determination of such domains from sequence information.

The invention also provides species homologs of the disclosed polynucleotides and proteins. Species homologs can be isolated and identified, for example, by making suitable probes or primers from the sequences disclosed herein and screening a suitable nucleic acid source from the desired species. The invention also encompasses allelic variants of the disclosed polynucleotides or proteins. Allelic variants are naturally-occurring alternative forms of polynucleotides which encode proteins which are identical, homologous, or related to those encoded by the polynucleotides shown in SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, and 43.

Proteins of the invention can be prepared by culturing transformed host cells under culture conditions suitable for expression of the recombinant protein. If a protein of the invention is produced in a yeast or bacterial expression system, it may be necessary to modify the protein, for example, by phosphorylation or glycosylation of appropriate sites, in order to obtain the protein in a functional form. Such covalent

attachments can be made using known chemical or enzymatic methods. The resulting expressed protein can then be purified from the culture (*i.e.*, from culture medium or cell extracts) using known purification techniques, such as size exclusion chromatography, ammonium sulfate fractionation, ion exchange chromatography, affinity  
5 chromatography, crystallization, electrofocusing, immunoprecipitation, immunoaffinity chromatography, and preparative gel electrophoresis.

A protein of the invention can optionally be expressed in a form which will facilitate purification. A protein can be expressed as a fusion protein with, for example, maltose binding protein (MBP), glutathione-S-transferase (GST), or thioredoxin (TRX).  
10 Kits for expression and purification of such fusion proteins are commercially available from New England BioLab (Beverly, Mass.), Pharmacia (Piscataway, N.J.) and In Vitrogen, respectively. Alternatively, a protein of the invention can be tagged with an epitope and subsequently purified using a specific antibody directed to the epitope. One such epitope, Flag, is commercially available from Kodak (New Haven, Conn.).

15 A protein of the invention can be expressed as a product of transgenic animals, *e.g.*, as a component of the milk of transgenic cows, goats, pigs, or sheep which are characterized by somatic or germ cells containing a nucleotide sequence encoding the protein. Proteins of the invention can also be produced by known conventional chemical synthesis. Methods for constructing the proteins of the present invention by synthetic  
20 means, such as solid phase peptide synthesis, are well known in the art.

Fusion proteins comprising amino acid sequences of proteins of the invention can also be constructed. Fusion proteins are useful for generating antibodies against amino acid sequences and for use in various assay systems. For example, fusion proteins can be used to identify proteins which interact with proteins of the invention. Physical methods,  
25 such as protein affinity chromatography, or library-based assays for protein-protein interactions such as the yeast two-hybrid or phage display systems, can also be used for this purpose. Such methods are well known in the art and can also be used as drug screens.

A fusion protein of the invention comprises two protein segments fused together  
30 by means of a peptide bond. The first protein segment consists of at least 95, 100, 120,



130, or 140 contiguous amino acids of SEQ ID NO:2, at least 6, 8, 10, 20, 30, 40, 50, 60, 70, 80, 90, 100, 101, 110, 120, 130, 150, 160, 170, or 180 contiguous amino acids of SEQ ID NO:4, at least 14, 15, 16, 18, 20, 25, or 30 contiguous amino acids selected from amino acids 1-312 of SEQ ID NO:6 or at least 75, 100, 125, 150, 175, 179, 200, 225, 250, 275, 300, 325, or 350 contiguous amino acids of SEQ ID NO:6, at least 17, 18, 19, 20, 25, or 30 contiguous amino acids selected from amino acids 1-287 of SEQ ID NO:8 or at least 136, 140, 150, 150, 179, 200, 250, 300, 350, or 400 contiguous amino acids selected from SEQ ID NO:8, at least 31, 32, 35, 40, or 45 contiguous amino acids selected from amino acids 1-238 of SEQ ID NO:10, or at least 82, 85, 100, 132, 150, 200, 225, 250, or 275 contiguous amino acids of SEQ ID NO:10, at least 6, 7, 8, 9, 10, 15, or 20 contiguous amino acids selected from amino acids 1-184 or 270-362 of SEQ ID NO: 12, at least 8, 9, 10, 12, 15, 20, or 25 contiguous amino acids selected from amino acids 268-364 of SEQ ID NO: 12, at least 27, 30, 35, or 40 contiguous amino acids selected from amino acids 250-383 of SEQ ID NO:12, or at least 96, 100, 150, 200, 250, 300, or 350 contiguous amino acids selected from SEQ ID NO:12, at least 6, 7, 8, 9, 10, 12, 15, or 20 contiguous amino acids selected from amino acids 1-111 or 204-261 of SEQ ID NO: 14, at least 17, 18, 20, 25, or 30 contiguous amino acids selected from amino acids 1-150 of SEQ ID NO:14, at least 75, 80, 100, 104, 125, 150, 175, 200, 225, or 250 contiguous amino acids of SEQ ID NO:14, at least 8, 10, 12, 14, 16, 20, 30, 40, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, 500, 525, or 550 contiguous amino acids of SEQ ID NO:16, at least 39, 40, 45, 46, or 50 contiguous amino acids selected from amino acids 13-232 of SEQ ID NO:18 or at least 46, 50, 55, 60, 75, 100, 125, 150, 175, 200, or 225 contiguous amino acids of SEQ ID NO:18, at least 6, 8, 10, 12, 15, 20, 25, 30, 50, 75, 100, 125, or 140 contiguous amino acids from SEQ ID NO:20, at least 7, 8, 10, 12, 15, 20, 25, 30, 50, 75, 100, 125, 150, or 160 contiguous amino acids from SEQ ID NO:22, at least 7, 8, 10, 12, 15, 20, 25, 30, 50, 75, 100, 125, 150, 175, 200, 225, 250, or 275 contiguous amino acids of SEQ ID NO:24, at least 11, 12, 15, 18, 20, 25, 30, 35, 50, 75, 100, 125, 150, 175, 200, or 225 contiguous amino acids of SEQ ID NO:26, at least 6, 8, 10, 12, 15, or 20 contiguous amino acids selected from amino acids 1-31 of SEQ ID NO:28 or at least 257, 260, 270, 280, 290,

300, 325, 350, 375, 400, 425, or 450 contiguous amino acids of SEQ ID NO:28, at least 6, 8, 10, 12, 15, 18, 20, 25, 30, 50, 75, 100, 125, 150, 175, or 200 contiguous amino acids of SEQ ID NO:30, at least 6, 8, 10, 12, 15, or 20 contiguous amino acids selected from amino acids 1-65 of SEQ ID NO:32 or at least 117, 120, 150, 175, 200, or 225  
5 contiguous amino acids of SEQ ID NO:32, at least 6, 8, 10, 12, 15, 20, 25, 30, 50, 75, 100, 125, 150, or 175 contiguous amino acids of SEQ ID NO:34, at least 14, 15, 18, 20, 25, 30, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, or 300 contiguous amino acids of SEQ ID NO:36, at least 19, 20, 25, 30, 35, 40, 50, 75, 100, 125, 150, 175, 200, 224, 250, 275, 300, or 325 contiguous amino acids of SEQ ID NO:38, at least 8, 10, 12, 15, 18, 20,  
10 25, 30, 50, 75, 100, 125, 150, 175, 200, 225, 250, 275, 300, 325, 350, 375, 400, 425, 450, 475, or 500 contiguous amino acids of SEQ ID NO:40, at least 7, 8, 10, 12, 15, 20, 30, 50, 75, 100, or 125 contiguous amino acids of SEQ ID NO:42, at least 10, 12, 15, 20, 25, 30, 50, 75, 100, 125, 150, or 170 contiguous amino acids of SEQ ID NO:44. The amino acids can also be selected from biologically active variants of those sequences. The first  
15 protein segment can also be a full-length protein as shown in SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, or 44. The first protein segment can be N-terminal or C-terminal, as is convenient.

The second protein segment can be a full-length protein or a protein fragment or polypeptide. Proteins commonly used in fusion protein construction include  $\beta$ -  
20 galactosidase,  $\beta$ -glucuronidase, green fluorescent protein (GFP), autofluorescent proteins, including blue fluorescent protein (BFP), glutathione-S-transferase (GST), luciferase, horseradish peroxidase (HRP), and chloramphenicol acetyltransferase (CAT). Epitope tags can be used in fusion protein constructions, including histidine (His) tags, FLAG tags, influenza hemagglutinin (HA) tags, Myc tags, VSV-G tags, and thioredoxin  
25 (Trx) tags. Other fusion constructions can include maltose binding protein (MBP), S-tag, Lex A DNA binding domain (DBD) fusions, GAL4 DNA binding domain fusions, and herpes simplex virus (HSV) BP16 protein fusions.

Fusion proteins of the invention can be made by covalently linking the first and second protein segments or by standard procedures in the art of molecular biology.  
30 Recombinant DNA methods can be used to prepare fusion proteins, for example, by

making a DNA construct which comprises coding sequences selected from SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, or 43 in proper reading frame with nucleotides encoding the second protein segment and expressing the DNA construct in a host cell, as is known in the art. Many kits for constructing fusion proteins are available from companies which supply research labs with tools for experiments, including, for example, Promega Corporation (Madison, WI), Stratagene (La Jolla, CA), Clontech (Mountain View, CA), Santa Cruz Biotechnology (Santa Cruz, CA), MBL International Corporation (MIC; Watertown, MA), and Quantum Biotechnologies (Montreal, Canada; 1-888-DNA-KITS).

Isolated proteins, polypeptides, biologically active variants, or fusion proteins can be used as immunogens, to obtain a preparation of antibodies which specifically bind to epitopes of the secreted proteins disclosed herein. The entire protein or fragments of the protein can be used as an immunogen, optionally conjugated to a hapten, such as keyhole limpet hemocyanin.

The antibodies can be used, *inter alia*, to detect proteins of the invention in human tissue or in fractions thereof. The antibodies can also be used to detect the presence of mutations in the genes encoding these proteins which result in under- or over-expression of proteins of the invention or in expression of a secreted protein with altered size or electrophoretic mobility. By binding to a protein of the invention, antibodies can also alter the functions of the protein.

Antibodies which specifically bind to a protein of the invention can be useful diagnostic agents. Antibodies can also be used to treat conditions associated with the protein, including forms of cancer in which abnormal expression of the protein is involved. In the case of neoplastic cells, antibodies which specifically bind to the protein can be useful for suppressing the metastatic spread of the neoplastic cells, which can be mediated by the protein.

Antibodies which specifically bind to epitopes of the secreted proteins, polypeptides, fusion proteins, or biologically active variants disclosed herein can be used in immunochemical assays, including but not limited to Western blots, ELISAs, radioimmunoassays, immunohistochemical assays, immunoprecipitations, or other

immunochemical assays known in the art. Typically, antibodies of the invention provide a detection signal at least 5-, 10-, or 20-fold higher than a detection signal provided with other proteins when used in such immunochemical assays. Preferably, antibodies which specifically bind to epitopes of a particular secreted protein do not detect other proteins in immunochemical assays and can immunoprecipitate that protein or polypeptide fragments of the protein from solution.

Specific antibodies specifically bind to epitopes present in a secreted protein having one of the amino acid sequences shown in SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, or 44 or to biologically active variants of those sequences. Typically, at least 6, 8, 10, or 12 contiguous amino acids are required to form an epitope. However, epitopes which involve non-contiguous amino acids may require more, *e.g.*, at least 15, 25, or 50 amino acids. Preferably, the epitopes are not present in other human proteins.

Epitopes of proteins of the invention which are particularly antigenic can be selected, for example, by routine screening of polypeptide fragments of the protein for antigenicity or by applying a theoretical method for selecting antigenic regions of a protein to the amino acid sequences disclosed herein. Such methods are taught, for example, in Hopp and Wood, *Proc. Natl. Acad. Sci. U.S.A.* 78, 3824-28 (1981), Hopp and Wood, *Mol. Immunol.* 20, 483-89 (1983), and Sutcliffe *et al.*, *Science* 219, 660-66 (1983).

Any type of antibody known in the art can be generated to bind specifically to epitopes of a secreted protein of the invention. For example, preparations of polyclonal and monoclonal antibodies can be made using standard methods which are well known in the art. Similarly, single-chain antibodies can also be prepared. Single-chain antibodies can be isolated, for example, from single-chain immunoglobulin display libraries, as is known in the art. The library is "panned" against amino acid sequences of a particular protein of the invention, and a number of single chain antibodies which bind with high-affinity to different epitopes of the protein can be isolated. Hayashi *et al.*, 1995, *Gene* 160:129-30. Single-chain antibodies can also be constructed using a DNA amplification method, such as the polymerase chain reaction (PCR), using hybridoma

cDNA as a template. Thirion *et al.*, 1996, *Eur. J. Cancer Prev.* 5:507-11.

Single-chain antibodies can be mono- or bispecific, and can be bivalent or tetravalent. Construction of tetravalent, bispecific single-chain antibodies is taught, for example, in Coloma and Morrison, 1997, *Nat. Biotechnol.* 15:159-63. Construction of  
5 bivalent, bispecific single-chain antibodies is taught *inter alia* in Mallender and Voss, 1994, *J. Biol. Chem.* 269:199-206.

A nucleotide sequence encoding a single-chain antibody can be constructed using manual or automated nucleotide synthesis, cloned into an expression construct using standard recombinant DNA methods, and introduced into a cell to express the coding  
10 sequence, as described below. Alternatively, single-chain antibodies can be produced directly using, for example, filamentous phage technology. Verhaar *et al.*, 1995, *Int. J. Cancer* 61:497-501; Nicholls *et al.*, 1993, *J. Immunol. Meth.* 165:81-91.

Monoclonal and other antibodies can also be "humanized" in order to prevent a patient from mounting an immune response against the antibody when it is used  
15 therapeutically. Such antibodies may be sufficiently similar in sequence to human antibodies to be used directly in therapy or may require alteration of a few key residues. Sequence differences between, for example, rodent antibodies and human sequences can be minimized by replacing residues which differ from those in the human sequences, for example, by site directed mutagenesis of individual residues, or by grafting of entire  
20 complementarity determining regions. Alternatively, one can produce humanized antibodies using recombinant methods, as described in GB2188638B. Antibodies which specifically bind to epitopes of a protein of the invention can contain antigen binding sites which are either partially or fully humanized, as disclosed in U.S. 5,565,332.

Other types of antibodies can be constructed and used in methods of the  
25 invention. For example, chimeric antibodies can be constructed as disclosed, for example, in WO 93/03151. Binding proteins which are derived from immunoglobulins and which are multivalent and multispecific, such as the "diabodies" described in WO 94/13804, can also be prepared.

Antibodies of the invention can be purified by methods well known in the art.  
30 For example, antibodies can be affinity purified by passing the antibodies over a column

to which a protein, polypeptide, biologically active variant, or fusion protein of the invention is bound. The bound antibodies can then be eluted from the column, using a buffer with a high salt concentration.

Specific-binding polypeptides other than antibodies can also be generated.

5 Specific-binding polypeptides are polypeptides which bind with a secreted protein or its variants and which have a measurably higher binding affinity for that protein and polypeptide fragments or variants of the protein than for other polypeptides tested for binding. Higher affinity by a factor of 10 is preferred, more preferably a factor of 100. Such polypeptides can be found, for example, using the yeast two-hybrid system.

10 Polynucleotides and proteins of the present invention exhibit one or more of the utilities or biological activities which are identified below. Biological activities and utilities of proteins of the invention can be provided by administration or use of the proteins themselves or by administration or use of polynucleotides encoding the proteins.

A protein of the invention can exhibit cytokine, cell proliferation (either inducing  
15 or inhibiting), or cell differentiation (either inducing or inhibiting) activity, or can induce production of other cytokines in certain cell populations. Many protein factors discovered to date, including all known cytokines, have exhibited activity in one or more factor-dependent cell proliferation assays; hence, the assays serve as a convenient confirmation of cytokine activity. The activity of a protein of the invention can be  
20 evidenced by any one of a number of routine factor-dependent cell proliferation assays for cell lines including, 32D (a mouse IL-3-dependent lymphoblast cell line, ATCC No. CRL-11346), DA2, DA1G, T10 (a human myeloma cell line, ATCC No. CRL-9068), B9, B9/11, BaF3, MC9/G, M+(preB M+), 2E8 (a mouse IL-7-dependent lymphoblast cell line, ATCC No. TIB-239), RB5, DA1, 123, T1165, HT2 (a mouse lymphoma cell  
25 line, ATCC No. CRL-8629), CTLL2, TF-1 (a human IL-5-unresponsive lymphoblast cell line, ATCC No. CRL-2003), Mo7e, and CMK.

Assays for T-cell or thymocyte proliferation include those described in CURRENT  
PROTOCOLS IN IMMUNOLOGY, Coligan *et al.*, eds., Greene Publishing Associates and  
Wiley-Interscience (particularly chapter 3, *In Vitro* Assays for Mouse Lymphocyte  
30 Function 3.1-3.19; and chapter 7, Immunologic Studies in Humans); Takai *et al.*, *J.*

*Immunol.* 137:3494-3500, 1986; Bertagnolli *et al.*, *J. Immunol.* 145:1706-1712, 1990; Bertagnolli *et al.*, *Cellular Immunology* 133:327-341, 1991; Bertagnolli, *et al.*, *J. Immunol.* 149:3778-3783, 1992; and Bowman *et al.*, *J. Immunol.* 152:1756-1761, 1994.

5 Assays for cytokine production and/or proliferation of spleen cells, lymph node cells, or thymocytes include those described in Kruisbeek and Shevach, *Polyclonal T Cell Stimulation*, in CURRENT PROTOCOLS IN IMMUNOLOGY, vol. 1, pp. 3.12.1-3.12.14, and Schreiber, *Measurement of Mouse and Human Interleukin Gamma*, in CURRENT PROTOCOLS IN IMMUNOLOGY, vol. 1, pp. 6.8.1-6.8.8.

10 Assays for proliferation and differentiation of hematopoietic and lymphopoietic cells include those described in Bottomly, *Measurement of Human and Murine Interleukin 2 and Interleukin 4*, in CURRENT PROTOCOLS IN IMMUNOLOGY, vol. 1, pp. 6.3.1-6.3.12; deVries *et al.*, *J. Exp. Med.* 173:1205-1211, 1991; Moreau *et al.*, *Nature* 336:690-692, 1988; Greenberger *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 80:2931-2938, 1983; Nordan, R., *Measurement of mouse and human interleukin 6*, in CURRENT  
15 PROTOCOLS IN IMMUNOLOGY, vol. 1, pp. 6.6.1-6.6.5; Smith *et al.*, *Proc. Natl. Acad. Sci. U.S.A.* 83:1857-1861, 1986; Bennett *et al.*, *Measurement of Human Interleukin 11*, in CURRENT PROTOCOLS IN IMMUNOLOGY, vol. 1, pp. 6.15.1; Ciarletta *et al.*, *Measurement of mouse and human Interleukin 9*, in CURRENT PROTOCOLS IN IMMUNOLOGY, vol. 1, p. 6.13.1.

20 Assays for T cell clone responses to antigens (which will identify, among others, proteins that affect APC-T cell interactions as well as direct T cell effects by measuring proliferation and cytokine production) include those described in CURRENT PROTOCOLS IN IMMUNOLOGY, especially chapters 3 (*In Vitro* Assays for Mouse Lymphocyte Function), chapter 6 (Cytokines and Their Cellular Receptors), and chapter 7  
25 (Immunologic Studies in Humans); Weinberger *et al.*, *Proc. Natl. Acad. Sci. USA* 77:6091-6095, 1980; Weinberger *et al.*, *Eur. J. Immun.* 11:405-411, 1981; Takai *et al.*, *J. Immunol.* 137:3494-3500, 1986; and Takai *et al.*, *J. Immunol.* 140:508-512, 1988.

A protein of the present invention can be useful to support colony forming cells or factor-dependent cell lines, to regulate hematopoiesis, and to treat myeloid or lymphoid cell deficiencies. Such proteins can be used, either alone or in combination with other cytokines, to support the growth and proliferation of erythroid progenitor  
5 cells. The proteins can also be used to treat various anemias, in conjunction with irradiation or chemotherapy to stimulate the production of erythroid precursors or erythroid cells.

A protein of the invention can have CSF activity and can be used to support the growth and proliferation of myeloid cells, such as granulocytes, monocytes, or  
10 macrophages. Proteins with such activity can be used, for example, in conjunction with chemotherapy to prevent or treat myelo-suppression. Proteins of the invention can also be used to support the growth and proliferation of megakaryocytes and platelets, thereby allowing prevention or treatment of platelet disorders such as thrombocytopenia. Proteins with such activity can be used to support the growth and proliferation of  
15 hematopoietic stem cells, either in place of or in conjunction with platelet transfusions. Proteins of the invention can be used to treat stem cell disorders, such as aplastic anemia and paroxysmal nocturnal hemoglobinuria, or to repopulate the stem cell compartment after irradiation or chemotherapy, either *in-vivo* or *ex-vivo*. For example, a protein of the invention can be used in conjunction with homologous or heterologous bone marrow  
20 transplantation or peripheral progenitor cell transplantation.

Suitable assays for proliferation and differentiation of various hematopoietic lines are cited above. Assays for embryonic stem cell differentiation which can identify proteins which influence embryonic hematopoiesis include those described in Johansson  
25 *et al. Cellular Biology* 15:141-151, 1995; Keller *et al., Molecular and Cellular Biology* 13:473-486, 1993; and McClanahan *et al., Blood* 81:2903-2915, 1993.

Assays for stem cell survival and differentiation include those described in Freshney, *Methylcellulose colony forming assays*, in CULTURE OF HEMATOPOIETIC CELLS, Freshney *et al.* eds., pp. 265-268, Wiley-Liss, Inc., New York, N.Y. 1994; Hirayama *et al., Proc. Natl. Acad. Sci. USA* 89:5907-5911, 1992; McNiece and Briddell,  
30 *Primitive hematopoietic colony forming cells with high proliferative potential*, in



- CULTURE OF HEMATOPOIETIC CELLS, pp. 23-39; Neben *et al.*, *Experimental Hematology* 22:353-359, 1994; Ploemacher, *Cobblestone area forming cell assay*, in CULTURE OF HEMATOPOIETIC CELLS, pp. 1-21; Spooncer *et al.*, *Long term bone marrow cultures in the presence of stromal cells*, in CULTURE OF HEMATOPOIETIC CELLS, pp. 163-179;
- 5 Sutherland, *Long term culture initiating cell assay*, in CULTURE OF HEMATOPOIETIC CELLS, pp. 139-162. Such assays can be used to identify proteins which regulate lympho-hematopoiesis.

Compositions of the invention relate to isolated (purified) polypeptides and polynucleotides. These compositions are substantially free of other human proteins or

10 human polynucleotides. A composition containing A is "substantially free of" B when at least 85% by weight of the total A+B in the composition is A. Preferably, A comprises at least about 90% by weight of the total of A+B in the composition, more preferably at least about 96% or even 99% by weight.

A protein of the invention also can have utility in compositions used for growth

15 or differentiation of bone, cartilage, tendon, ligament, or nerve tissue, as well as for wound healing and tissue repair and replacement, and in the treatment of burns, incisions, and ulcers.

Proteins of the present invention can induce cartilage and/or bone growth in circumstances where bone is not normally formed and thus have an application in healing

20 bone fractures and cartilage damage or defects in humans and other animals. A preparation employing a protein of the invention can have prophylactic use in closed as well as open fracture reduction and also in the improved fixation of artificial joints. *De novo* bone formation induced by an osteogenic agent contributes to the repair of congenital, trauma- or surgery-induced craniofacial defects and also is useful in cosmetic

25 plastic surgery.

A protein of this invention can also be used in the treatment of periodontal disease and in other tooth repair processes. Such agents can provide an environment to attract bone-forming cells, stimulate growth of bone-forming cells, or induce differentiation of progenitors of bone-forming cells. A protein of the invention can be

30 used to treat osteoporosis or osteoarthritis, for example, through stimulation of bone

and/or cartilage repair or by blocking inflammation. Mechanisms of destroying tissue mediated by inflammatory processes, such as collagenase or osteoclast activity, can also be inhibited.

Tendon or ligament formation can also be influenced by a protein of the invention. A protein of the invention which induces tendon/ligament-like tissue or other tissue formation in circumstances where such tissue is not normally formed can be used to heal tendon or ligament tears, deformities, and other tendon or ligament defects in humans and other animals. A preparation employing a tendon/ligament-like tissue inducing protein can be used to prevent damage to tendon or ligament tissue, as well as in the improved fixation of tendon or ligament to bone or other tissues, and to repair defects to tendon or ligament tissue. *De novo* tendon/ligament-like tissue formation induced by a composition of the invention contributes to the repair of congenital, trauma-induced, or other tendon or ligament defects of other origin and can also be used in cosmetic plastic surgery, for attachment or repair of tendons or ligaments.

Compositions of the invention can provide an environment which will attract tendon- or ligament-forming cells, stimulate growth of tendon- or ligament-forming cells, induce differentiation of progenitors of tendon- or ligament-forming cells, or induce growth of tendon/ligament cells or progenitors *ex vivo*. Such cells can then be returned to the body to effect tissue repair. Compositions of the invention can also be used to treat tendinitis, carpal tunnel syndrome, and other tendon or ligament defects. Such compositions can optionally include an appropriate matrix and/or sequestering agent as a pharmaceutically acceptable carrier, as is well known in the art.

A protein of the invention can also be useful for proliferation of neural cells and for regeneration of nerve and brain tissue, *i.e.* for the treatment of central and peripheral nervous system diseases and neuropathies, as well as mechanical and traumatic disorders.

More specifically, a protein can be used in the treatment of diseases such as Alzheimer's disease, Parkinson's disease, Huntington's disease, amyotrophic lateral sclerosis, and Shy-Drager syndrome. Other conditions which can be treated in accordance with the invention include mechanical and traumatic disorders, such as spinal cord disorders and head trauma, and cerebrovascular diseases, such as stroke. Peripheral neuropathies

resulting from chemotherapy or other medical therapies can be treated using a protein of the invention.

Proteins of the invention can also be used to promote better or faster closure of non-healing wounds, including pressure ulcers, ulcers associated with vascular  
5 insufficiency, or surgical and traumatic wounds.

A protein of the invention can also affect generation or regeneration of other tissues, such as organs (including, for example, pancreas, liver, intestine, kidney, skin, endothelium), muscle (smooth, skeletal, or cardiac), and vascular (including vascular endothelium) tissue, or for promoting the growth of cells of which such tissues are  
10 comprised. Part of the desired effects can be by inhibition or modulation of fibrotic scarring to allow normal tissue to regenerate. A protein of the invention can also exhibit angiogenic activity.

A protein of the present invention can be useful for gut protection or regeneration, and for treatment of lung or liver fibrosis, reperfusion injury in various tissues, and  
15 conditions resulting from systemic cytokine damage. A protein of the invention can also be useful for promoting or inhibiting differentiation of tissues described above from precursor tissues or cells or for inhibiting the growth of tissues described above.

Assays for tissue generation activity include those described for bone, cartilage, and tendon in WO 95/16035, for neuronal tissue in WO 95/05846, and for skin and  
20 endothelial tissue in WO 91/07491. Assays for wound healing activity include, for example, those described in Winter, EPIDERMAL WOUND HEALING, polypeptides 71-112 (Maibach and Rovee, eds.), Year Book Medical Publishers, Inc., Chicago, and Eaglstein and Mertz, *J. Invest. Dermatol* 71:382-84 (1978).

A protein of the present invention can also demonstrate activity as a receptor,  
25 receptor ligand, or inhibitor or agonist of a receptor/ligand interaction. Examples of such receptors and ligands include cytokine receptors and their ligands, receptor kinases and their ligands, receptor phosphatases and their ligands, receptors involved in cell-cell interactions and their ligands, including cellular adhesion molecules such as selectins, integrins, and their ligands, and receptor/ligand pairs involved in antigen presentation,  
30 antigen recognition and development of cellular and humoral immune responses.

Receptors and ligands are also useful for screening of potential peptide or small molecule inhibitors of the relevant receptor/ligand interaction. A protein of the invention, including fragments of receptors and ligands, can itself be useful as an inhibitor of receptor/ligand interactions.

- 5           Suitable assays for receptor-ligand activity include those described in CURRENT PROTOCOLS IN IMMUNOLOGY, chapter 7.28, *Measurement of Cellular Adhesion under static conditions*, pages 7.28.1-7.28.22, Takai *et al.*, *Proc. Natl. Acad. Sci. USA* 84:6864-6868, 1987; Bierer *et al.*, *J. Exp. Med.* 168:1145-1156, 1988; Rosenstein *et al.*, *J. Exp. Med.* 169:149-160 1989; Stoltenborg *et al.*, *J. Immunol. Methods* 175:59-68, 10   1994; Stitt *et al.*, *Cell* 80:661-670, 1995.

- A protein of the invention can be used in a pharmaceutical composition. Compositions comprising proteins or polynucleotides of the invention have therapeutic applications, both for human patients and veterinary patients, such as domestic animals and thoroughbred horses. Such compositions can optionally include a pharmaceutically 15   acceptable carrier. In addition to protein and carrier, such a composition can also contain diluents, fillers, salts, buffers, stabilizers, solubilizers, and other materials well known in the art. Characteristics of a carrier will depend on the route of administration. Compositions of the invention can also contain cytokines, lymphokines, or other hematopoietic factors such as M-CSF, GM-CSF, TNF, IL-1, IL-2, IL-3, IL-4, IL-5, IL-6, 20   IL-7, IL-8, IL-9, IL-10, IL-11, IL-12, IL-13, IL-14, IL-15, IFN, TNF0, TNF1, TNF2, G-CSF, Meg-CSF, thrombopoietin, stem cell factor, erythropoietin, or growth factors such as epidermal growth factor (EGF), platelet derived growth factor (PDGF), transforming growth factors (TGF- $\alpha$  and TGF- $\beta$ ), or insulin-like growth factor (IGF).

- A pharmaceutical composition can also contain other agents which either enhance 25   the activity of the protein or complement its activity or use in treatment. Such additional factors and/or agents can be included in the pharmaceutical composition to produce a synergistic effect with a protein of the invention or to minimize side effects. Conversely, a protein of the invention can be included in formulations of a particular factor, such as a cytokine, lymphokine, other hematopoietic factor, thrombolytic or anti-thrombotic factor, 30   or anti-inflammatory agent to minimize side effects of the factor.

A protein of the present invention can be active in multimers (*e.g.*, heterodimers or homodimers) or complexes with itself or other proteins, and compositions of the invention can comprise a protein of the invention in such a multimeric or complexed form. For example, a composition of the invention can be in the form of a complex of a protein or proteins of the invention together with protein or peptide antigens. The protein or peptide antigen will deliver a stimulatory signal to both B and T lymphocytes. B lymphocytes will respond to antigen through their surface immunoglobulin receptor. T lymphocytes will respond to antigen through the T cell receptor (TCR) following presentation of the antigen by MHC proteins. MHC proteins and structurally related proteins, including those encoded by class I and class II MHC genes on host cells, can present the peptide antigen(s) to T lymphocytes. Antigen components can also be supplied as purified MHC-peptide complexes alone or with co-stimulatory molecules which can directly signal T cells. Alternatively, antibodies able to bind surface immunoglobulin and other molecules on B cells, as well as antibodies able to bind the TCR and other molecules on T cells, can be combined with a composition of the invention.

A composition of the invention can be in the form of a liposome in which a protein of the invention is combined, in addition to other pharmaceutically acceptable carriers, with amphipathic agents such as lipids, which exist in aggregated form as micelles, insoluble monolayers, liquid crystals, or lamellar layers in aqueous solution. Suitable lipids for liposomal formulation include monoglycerides, diglycerides, sulfatides, lysolecithin, phospholipids, saponin, bile acids, and the like. Preparation of liposomal formulations is within the level of skill in the art, as disclosed, for example, in U.S. 4,235,871, U.S. 4,501,728, U.S. 4,837,028, and U.S. 4,737,323.

A therapeutically effective amount of a protein of the invention is administered to a mammal having a condition to be treated. The amount of protein which is therapeutically effective is that amount of protein which is sufficient to treat, heal, prevent, or ameliorate the condition, or to increase the rate of such treatment. Proteins of the invention can be administered either alone or in combination with other therapeutic agents, such as cytokines, lymphokines, or other hematopoietic factors. Other

therapeutic agents can be administered simultaneously or sequentially with proteins of the invention, as determined by the attending physician.

Compositions of the invention can be inhaled, ingested, applied topically, or administered by cutaneous, subcutaneous, intraperitoneal, parenteral or intravenous injection. When a therapeutically effective amount of protein of the present invention is administered orally, protein of the present invention will be in the form of a tablet, capsule, powder, solution or elixir. When administered in tablet form, the pharmaceutical composition of the invention can additionally contain a solid carrier such as a gelatin or an adjuvant. The tablet, capsule, and powder contain from about 5-95%, 25-90%, 30-80%, 40-75%, or 50% protein of the invention by weight. When administered in liquid form, a liquid carrier such as water, petroleum, oils of animal or plant origin such as peanut oil, mineral oil, soybean oil, or sesame oil, or synthetic oils can be added. The liquid form of the composition can further contain physiological saline solution, dextrose or other saccharide solution, or glycols such as ethylene glycol, propylene glycol, or polyethylene glycol. When administered in liquid form, the pharmaceutical composition contains from about 0.5-90%, 1-80%, 5-75%, 10-65%, 20-50%, 10-50%, or 25-40% by weight of protein of the invention.

When a therapeutically effective amount of protein of the present invention is administered by intravenous, cutaneous, or subcutaneous injection, a pyrogen-free, parenterally acceptable aqueous solution of the protein is preferred. The skilled artisan can readily prepare an acceptable protein solution with suitable pH, isotonicity, and stability. A solution of the composition for intravenous, cutaneous, or subcutaneous injection should also contain an isotonic vehicle, such as Sodium Chloride Injection, Ringer's Injection, Dextrose Injection, Dextrose and Sodium Chloride Injection, Lactated Ringer's Injection, or other vehicle as known in the art. Stabilizers, preservatives, buffers, antioxidants, or other additives known to those of skill in the art can also be added to the composition.

The amount of protein of the present invention in the pharmaceutical composition of the present invention will depend upon the nature and severity of the condition being treated and on the nature of prior treatments which the patient has undergone.

Ultimately, the attending physician will decide the amount of protein of the present invention with which to treat each individual patient. Initially, the attending physician will administer low doses of protein of the present invention and observe the patient's response. Larger doses of protein of the present invention can be administered until the  
5 optimal therapeutic effect is obtained for the patient, and at that point the dosage is not increased further. It is contemplated that the various pharmaceutical compositions used to practice the method of the present invention should contain about 0.01  $\mu$ g to about 100 mg (preferably about 0.1  $\mu$ g to about 10 mg, more preferably about 0.1  $\mu$ g to about 1 mg) of protein of the present invention per kg body weight.

10 Duration of intravenous therapy using a composition of the invention will vary, depending on the severity of the disease being treated and the condition and potential idiosyncratic response of each individual patient. It is contemplated that the duration of each application of a composition of the invention will be in the range of 12 to 24 hours of continuous intravenous administration. Ultimately, the attending physician will  
15 decide on the appropriate duration of intravenous therapy.

A composition of the invention which is useful for bone, cartilage, tendon or ligament regeneration can be administered topically, systematically, or locally in an implant or device. Encapsulation or injection in a viscous form for delivery to the site of bone, cartilage or tissue damage is also possible. Topical administration can be suitable  
20 for wound healing and tissue repair. Optionally, therapeutic agents other than a protein of the invention can be included in the composition, as described above.

To affect bone or cartilage formation, a composition of the invention would include a matrix capable of delivering the composition to the site of bone or cartilage damage and for providing a structure for the developing bone and cartilage. Optimally,  
25 the matrix would be capable of resorption into the body. Matrices can be formed of materials presently in use for other implanted medical applications, the choice of material being based on biocompatibility, biodegradability, mechanical properties, cosmetic appearance, and interface properties. Suitable biodegradable matrix materials include chemically defined calcium sulfate, tricalciumphosphate, hydroxyapatite, polylactic acid,  
30 polyglycolic acid, polyanhydride, bone or dermal collagen, pure proteins, and

extracellular matrix components. Suitable nonbiodegradable and chemically defined matrix materials include sintered hydroxyapatite, bioglass, aluminates, or other ceramics. Individual matrix components can be modified, for example, to affect pore size, particle size, particle shape, and biodegradability. Combinations of materials can be used, as is known in the art.

Sequestering agents, such as carboxymethyl cellulose or an autologous blood clot, can be employed to prevent protein compositions from dissociating from the matrix. Sequestering agents include cellulosic materials such as alkylcelluloses (including hydroxyalkylcelluloses), including methylcellulose, ethylcellulose, hydroxyethylcellulose, hydroxypropylcellulose, hydroxypropyl-methylcellulose, and carboxymethylcellulose, the most preferred being cationic salts of carboxymethylcellulose (CMC). Other preferred sequestering agents include hyaluronic acid, sodium alginate, polyethylene glycol, polyoxyethylene oxide, carboxyvinyl polymer and polyvinyl alcohol. The amount of sequestering agent is based on total formulation weight, such as 0.5-20% or 1-10%, and should be an amount of sequestering agent which prevents desorption of the protein from the polymer matrix but which permits progenitor cells to infiltrate the matrix, so that the protein can assist the osteogenic activity of the progenitor cells.

The dosage regimen of a protein-containing pharmaceutical composition to be used in tissue regeneration will be determined by the attending physician considering various factors which modify the action of the proteins, *e.g.*, amount of tissue weight desired to be formed, the site of damage, the condition of the damaged tissue, the size of a wound, type of damaged tissue (*e.g.*, bone), the patient's age, sex, and diet, the severity of any infection, time of administration, and other clinical factors. The dosage can vary with the type of matrix used in the reconstitution and whether other therapeutic agents, such as growth factors, are included. Progress of the treatment can be monitored by periodic assessment of tissue/bone growth and/or repair, for example, using X-rays, histomorphometric determinations, or tetracycline labeling.

Polynucleotides of the invention can also be used for gene therapy. Polynucleotides can be introduced either *in vivo* or *ex vivo* into cells for expression in a



mammalian subject. Cells can be cultured *ex vivo* in the presence of proteins of the invention in order to produce a desired effect on or activity in such cells. Treated cells can then be introduced *in vivo* for therapeutic purposes, as is known in the art.

Polynucleotides of the invention can be administered by known methods of introducing  
5 polynucleotides into a cell or organism (including in the form of viral vectors or naked DNA).

Polynucleotides of the invention can also be delivered to subjects for the purpose of screening test compounds for those which are useful for enhancing transfer of polynucleotides of the invention to a cell or for enhancing subsequent biological effects  
10 of the polynucleotides within the cell. Such biological effects include hybridization to complementary mRNA and inhibition of its translation, expression of the polynucleotide to form mRNA and/or protein, and replication and integration of the polynucleotide.

Test compounds which can be screened include any substances, whether natural products or synthetic, which can be administered to the subject. Libraries or mixtures of  
15 compounds can be tested. The compounds or substances can be those for which a pharmaceutical effect is previously known or unknown. The compounds or substances can be delivered before, after, or concomitantly with the polynucleotides. They can be administered separately or in admixture with the polynucleotides.

Integration of delivered polynucleotides can be monitored by any means known  
20 in the art. For example, Southern blotting of the delivered polynucleotides can be performed. A change in the size of the fragments of the delivered polynucleotides indicates integration. Replication of the delivered polynucleotides can be monitored *inter alia* by detecting incorporation of labeled nucleotides combined with hybridization to a specific nucleotide probe. Expression of a polynucleotide of the invention can be  
25 monitored by detecting production of mRNA which hybridizes to the delivered polynucleotide or by detecting protein. Proteins of the invention can be detected immunologically. Thus, delivery of polynucleotides of the invention according to the present invention provides an excellent system for screening test compounds for their ability to enhance delivery, integration, hybridization, expression, replication or  
30 integration in an animal, preferably a mammal, more preferably a human.

Polynucleotides of the invention can be used for a variety of research purposes. Any or all of these research utilities are capable of being developed into reagent grade or kit format for commercialization as research products. For example, polynucleotides can be used to express recombinant protein for analysis, characterization, or therapeutic use.

5 Polynucleotides can also be used as markers for tissues in which the corresponding protein is preferentially expressed, either constitutively or at a particular stage of tissue differentiation or development or in disease states. Polynucleotides can also be used as molecular weight markers on Southern gels or, when labeled, for example, with a fluorescent tag or a radiolabel, polynucleotides can be used as chromosome markers, to

10 identify chromosomes for gene mapping. Potential genetic disorders can be identified by comparing the sequences of wild-type polynucleotides of the invention with endogenous nucleotide sequences in patients. Polynucleotides of the invention can also be used as probes for the discovery of novel, related DNA sequences, to derive PCR primers for genetic fingerprinting, as probes to "subtract-out" known sequences in the process of

15 discovering other novel polynucleotides, for selecting and making oligomers for attachment to a gene chip or other support, to raise anti-protein antibodies using DNA immunization techniques, and as antigens, to raise anti-DNA antibodies or to elicit another immune response.

Where the polynucleotide encodes a protein which binds or potentially binds to

20 another protein, such as in a receptor-ligand interaction, the polynucleotide can also be used in interaction trap assays, such as the yeast two-hybrid assay, to identify polynucleotides encoding the protein with which binding occurs or to identify inhibitors of the binding interaction, for example in drug screening assays.

Proteins of the invention can similarly be used in assays to determine biological activity, including use in a panel of multiple proteins for high-throughput screening, to raise antibodies or to elicit another immune response, as a reagent in assays designed to quantitatively determine levels of the protein (or its receptor) in biological fluids, as markers for tissues in which the protein is preferentially expressed (either constitutively or at a particular stage of tissue differentiation or development or in a disease state), and to identify related receptors or ligands. Where the protein binds or potentially binds to

another protein such as, for example, in a receptor-ligand interaction, the protein can be used to identify the other protein with which binding occurs or to identify inhibitors of the binding interaction. Proteins involved in these binding interactions can also be used to screen for peptide or small molecule inhibitors or agonists of the binding interaction.

5 Polynucleotides of the invention can also be used on polynucleotide arrays. Polynucleotide arrays provide a high throughput technique that can assay a large number of polynucleotide sequences in a sample. This technology can be used as a diagnostic tool and as a tool to test for differential expression of genes having the coding sequences disclosed herein.

10 To create arrays, single-stranded polynucleotide probes can be spotted onto a substrate in a two-dimensional matrix or array. The single-stranded polynucleotide probes can comprise at least 6, 7, 8, 9, 10, 11, 12, 13, 14, 15, 16, 17, 18, 19, 20, 25, or 30 or more contiguous nucleotides selected from the nucleotide sequences shown in SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, or 43.

15 The substrate can be any substrate to which polynucleotide probes can be attached, including but not limited to glass, nitrocellulose, silicon, and nylon. Polynucleotide probes can be bound to the substrate by either covalent bonds or by non-specific interactions, such as hydrophobic interactions. Techniques for constructing arrays and methods of using these arrays are described in EP No. 0 799 897; PCT No. WO 20 97/29212; PCT No. WO 97/27317; EP No. 0 785 280; PCT No. WO 97/02357; U.S. Pat. No. 5,593,839; U.S. Pat. No. 5,578,832; EP No. 0 728 520; U.S. Pat. No. 5,599,695; EP No. 0 721 016; U.S. Pat. No. 5,556,752; PCT No. WO 95/22058; and U.S. Pat. No. 5,631,734. Commercially available polynucleotide arrays, such as Affymetrix GeneChip<sup>®</sup>, can also be used. Use of the GeneChip<sup>®</sup> to detect gene expression is 25 described, for example, in Lockhart *et al.*, Nature Biotechnology 14:1675 (1996); Chee *et al.*, Science 274:610 (1996); Hacia *et al.*, Nature Genetics 14:441, 1996; and Kozal *et al.*, Nature Medicine 2:753, 1996.

Biological samples comprising single-stranded polynucleotides can be labeled and then hybridized to the probes. Detectable labels which can be used include but are 30 not limited to radiolabels, biotinylated labels, fluorophors, and chemiluminescent labels.

Double stranded polynucleotides, comprising the labeled sample polynucleotides bound to polynucleotide probes, can be detected once the unbound portion of the sample is washed away. Biological samples in which expression of genes comprising polynucleotides of the invention can be examined include samples of diseased and non-  
5 diseased tissues, samples of tissues suspected of being diseased (particularly tissues suspected of being neoplastic), samples of different cell types, samples of cells at different developmental stages, samples of tissues from different species, and the like.

The complete contents of all references cited in this disclosure are expressly incorporated herein by reference. While certain embodiments of the invention have been  
10 described with particularity herein, those of skill in the art will recognize that various modifications of the invention can be made. It is understood that such modifications and variations are included within the scope of the appended claims.

**WE CLAIM:**

1. An isolated and purified protein comprising an amino acid sequence which is at least 85% identical to an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, and 44, wherein percent identity is determined using a Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 1.
2. The isolated and purified protein of claim 1 wherein the amino acid sequence comprises an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, and 44
3. An isolated and purified protein comprising an amino acid sequence selected from the group consisting of at least 95 contiguous amino acids of SEQ ID NO:2, at least 101 contiguous amino acids of SEQ ID NO:4, at least 14 contiguous amino acids selected from amino acids 1-312 of SEQ ID NO:6, at least 179 contiguous amino acids of SEQ ID NO:6, at least 75 contiguous amino acids of SEQ ID NO:6, at least 179 contiguous amino acids of SEQ ID NO:8, at least 136 contiguous amino acids of SEQ ID NO:8, at least 17 contiguous amino acids selected from amino acids 1-287 of SEQ ID NO:8, at least 82 contiguous amino acids of SEQ ID NO:10, at least 31 contiguous amino acids selected from amino acids 1-238 of SEQ ID NO:10, at least 96 contiguous amino acids of SEQ ID NO:12, at least 27 contiguous amino acids selected from amino acids 250-383 of SEQ ID NO:12, at least 6 contiguous amino acids selected from amino acids 1-184 of SEQ ID NO:12, at least 8 contiguous amino acids selected from amino acids 268-364 of SEQ ID NO:12, at least 104 contiguous amino acids of SEQ ID NO:14, at least 75 contiguous amino acids of SEQ ID NO:14, at least 17

contiguous amino acids selected from amino acids 1-150 of SEQ ID NO:14, at least 6 contiguous amino acids selected from amino acids 204-261 of SEQ ID NO:14, at least 6 contiguous amino acids selected from amino acids 1-111 of SEQ ID NO:14, at least 8 contiguous amino acids of SEQ ID NO:16, at least 46 contiguous amino acids of SEQ ID NO:18, at least 39 contiguous amino acids selected from amino acids 13-232 of SEQ ID NO:18, at least 6 contiguous amino acids of SEQ ID NO:20, at least 7 contiguous amino acids of SEQ ID NO:22, at least 7 contiguous amino acids of SEQ ID NO:24, at least 11 contiguous amino acids of SEQ ID NO:26, at least 257 contiguous amino acids of SEQ ID NO:28, at least 6 contiguous amino acids selected from amino acids 1-31 of SEQ ID NO:28, at least 6 contiguous amino acids of SEQ ID NO:30, at least 117 contiguous amino acids of SEQ ID NO:32, at least 6 contiguous amino acids selected from amino acids 1-65 of SEQ ID NO:32, at least 6 contiguous amino acids of SEQ ID NO:34, at least 14 contiguous amino acids of SEQ ID NO:36, at least 19 contiguous amino acids of SEQ ID NO:38, at least 8 contiguous amino acids of SEQ ID NO:40, at least 7 contiguous amino acids of SEQ ID NO:42, and at least 10 contiguous amino acids of SEQ ID NO:44.

4. A fusion protein comprising two protein segments joined together with a peptide bond, wherein the first protein segment consists of an amino acid sequence selected from the group consisting of at least 95 contiguous amino acids of SEQ ID NO:2, at least 101 contiguous amino acids of SEQ ID NO:4, at least 14 contiguous amino acids selected from amino acids 1-312 of SEQ ID NO:6, at least 179 contiguous amino acids of SEQ ID NO:6, at least 75 contiguous amino acids of SEQ ID NO:6, at least 179 contiguous amino acids of SEQ ID NO:8, at least 136 contiguous amino acids of SEQ ID NO:8, at least 17 contiguous amino acids selected from amino acids 1-287 of SEQ ID NO:8, at least 82 contiguous amino acids of SEQ ID NO:10, at least 31 contiguous amino acids selected from amino acids 1-238 of SEQ ID NO:10, at least 96 contiguous amino acids of SEQ ID NO:12, at least 27 contiguous amino acids selected from amino acids 250-383

- of SEQ ID NO:12, at least 6 contiguous amino acids selected from amino acids 1-184 of SEQ ID NO:12, at least 8 contiguous amino acids selected from amino acids 268-364 of SEQ ID NO:12, at least 104 contiguous amino acids of SEQ ID NO:14, at least 75 contiguous amino acids of SEQ ID NO:14, at least 17 contiguous amino acids selected from amino acids 1-150 of SEQ ID NO:14, at least 6 contiguous amino acids selected from amino acids 204-261 of SEQ ID NO:14, at least 6 contiguous amino acids selected from amino acids 1-111 of SEQ ID NO:14, at least 8 contiguous amino acids of SEQ ID NO:16, at least 46 contiguous amino acids of SEQ ID NO:18, at least 39 contiguous amino acids selected from amino acids 13-232 of SEQ ID NO:18, at least 6 contiguous amino acids of SEQ ID NO:20, at least 7 contiguous amino acids of SEQ ID NO:22, at least 7 contiguous amino acids of SEQ ID NO:24, at least 11 contiguous amino acids of SEQ ID NO:26, at least 257 contiguous amino acids of SEQ ID NO:28, at least 6 contiguous amino acids selected from amino acids 1-31 of SEQ ID NO:28, at least 6 contiguous amino acids of SEQ ID NO:30, at least 117 contiguous amino acids of SEQ ID NO:32, at least 6 contiguous amino acids selected from amino acids 1-65 of SEQ ID NO:32, at least 6 contiguous amino acids of SEQ ID NO:34, at least 14 contiguous amino acids of SEQ ID NO:36, at least 19 contiguous amino acids of SEQ ID NO:38, at least 8 contiguous amino acids of SEQ ID NO:40, at least 7 contiguous amino acids of SEQ ID NO:42, and at least 10 contiguous amino acids of SEQ ID NO:44.
5. A preparation of antibodies which specifically binds to a protein comprising an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, and 44.
  6. An isolated and purified subgenomic polynucleotide which encodes a protein comprising an amino acid sequence which is at least 85% identical to an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, and 44, wherein

percent identity is determined using a Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 1.

7. The isolated and purified subgenomic polynucleotide of claim 6 wherein the amino acid sequence is selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, and 44.
8. An isolated and purified subgenomic polynucleotide comprising a nucleotide sequence which is at least 85% identical to a nucleotide sequence selected from the group consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 31, 33, 35, 37, 39, 41, 43, 45, and the complements thereof, wherein percent identity is determined using a Smith-Waterman homology search algorithm using an affine gap search with a gap open penalty of 12 and a gap extension penalty of 1.
9. An isolated and purified subgenomic polynucleotide which encodes an amino acid sequence selected from the group consisting of at least 95 contiguous amino acids of SEQ ID NO:2, at least 101 contiguous amino acids of SEQ ID NO:4, at least 14 contiguous amino acids selected from amino acids 1-312 of SEQ ID NO:6, at least 179 contiguous amino acids of SEQ ID NO:6, at least 75 contiguous amino acids of SEQ ID NO:6, at least 179 contiguous amino acids of SEQ ID NO:8, at least 136 contiguous amino acids of SEQ ID NO:8, at least 17 contiguous amino acids selected from amino acids 1-287 of SEQ ID NO:8, at least 82 contiguous amino acids of SEQ ID NO:10, at least 31 contiguous amino acids selected from amino acids 1-238 of SEQ ID NO:10, at least 96 contiguous amino acids of SEQ ID NO:12, at least 27 contiguous amino acids selected from amino acids 250-383 of SEQ ID NO:12, at least 6 contiguous amino acids selected from amino acids 1-184 of SEQ ID NO:12, at least 8 contiguous amino acids selected from amino acids 268-364 of SEQ ID NO:12, at least 104



contiguous amino acids of SEQ ID NO:14, at least 75 contiguous amino acids of SEQ ID NO:14, at least 17 contiguous amino acids selected from amino acids 1-150 of SEQ ID NO:14, at least 6 contiguous amino acids selected from amino acids 204-261 of SEQ ID NO:14, at least 6 contiguous amino acids selected from amino acids 1-111 of SEQ ID NO:14, at least 8 contiguous amino acids of SEQ ID NO:16, at least 46 contiguous amino acids of SEQ ID NO:18, at least 39 contiguous amino acids selected from amino acids 13-232 of SEQ ID NO:18, at least 6 contiguous amino acids of SEQ ID NO:20, at least 7 contiguous amino acids of SEQ ID NO:22, at least 7 contiguous amino acids of SEQ ID NO:24, at least 11 contiguous amino acids of SEQ ID NO:26, at least 257 contiguous amino acids of SEQ ID NO:28, at least 6 contiguous amino acids selected from amino acids 1-31 of SEQ ID NO:28, at least 6 contiguous amino acids of SEQ ID NO:30, at least 117 contiguous amino acids of SEQ ID NO:32, at least 6 contiguous amino acids selected from amino acids 1-65 of SEQ ID NO:32, at least 6 contiguous amino acids of SEQ ID NO:34, at least 14 contiguous amino acids of SEQ ID NO:36, at least 19 contiguous amino acids of SEQ ID NO:38, at least 8 contiguous amino acids of SEQ ID NO:40, at least 7 contiguous amino acids of SEQ ID NO:42, and at least 10 contiguous amino acids of SEQ ID NO:44.

10. The isolated and purified subgenomic polynucleotide of claim 9 which encodes an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, and 44.
11. The isolated and purified subgenomic polynucleotide of claim 10 wherein the nucleotide sequence is selected from the group consisting of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 31, 33, 35, 37, 39, 41, and 43.
12. An isolated and purified subgenomic polynucleotide comprising a polynucleotide segment which hybridizes to a nucleotide sequence selected from the group

consisting of SEQ ID NOS: 1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 31, 33, 35, 37, 39, 41, and 43, and the complements thereof after washing with 0.2X SSC at 65 °C, wherein the polynucleotide segment encodes a protein having an amino acid sequence selected from the group consisting of SEQ ID NOS:2, 4, 6, 8, 10, 12, 14, 16, 18, 20, 22, 24, 26, 28, 30, 32, 34, 36, 38, 40, 42, and 44.

13. An isolated and purified subgenomic polynucleotide comprising a nucleotide sequence selected from the group consisting of at least 499 contiguous nucleotides of SEQ ID NO:1, at least 1141 contiguous nucleotides of SEQ ID NO:1, at least 475 contiguous nucleotides of SEQ ID NO:3, at least 313 contiguous nucleotides selected from nucleotides 1-1001 of SEQ ID NO:3, at least 751 contiguous nucleotides of SEQ ID NO:5, at least 538 contiguous nucleotides of SEQ ID NO:5, at least 11 contiguous nucleotides selected from nucleotides 1-946 of SEQ ID NO:5, at least 13 contiguous nucleotides selected from nucleotides 1-1039 of SEQ ID NO:5, at least 651 contiguous nucleotides of SEQ ID NO:7, at least 522 contiguous nucleotides of SEQ ID NO:7, at least 11 contiguous nucleotides selected from nucleotides 1-913 of SEQ ID NO:7, at least 484 contiguous nucleotides of SEQ ID NO:9, at least 317 contiguous nucleotides of SEQ ID NO:9, at least 11 contiguous nucleotides selected from nucleotides 1-216 of SEQ ID NO:9, at least 11 contiguous nucleotides selected from nucleotides 379-812 of SEQ ID NO:9, at least 183 contiguous nucleotides selected from nucleotides 1-984 of SEQ ID NO:9, at least 594 contiguous nucleotides of SEQ ID NO:11, at least 289 contiguous nucleotides of SEQ ID NO:11, at least 11 contiguous nucleotides selected from nucleotides 1-585 of SEQ ID NO:11, at least 11 contiguous nucleotides selected from nucleotides 853-1120 of SEQ ID NO:11, at least 592 contiguous nucleotides of SEQ ID NO:13, at least 275 contiguous nucleotides of SEQ ID NO:13, at least 11 contiguous nucleotides selected from nucleotides 1-294 of SEQ ID NO:13, at least 537 contiguous nucleotides of SEQ ID NO:15, at least 294 contiguous nucleotides selected from nucleotides 1-1889 of SEQ ID NO:15, at least 171 contiguous

nucleotides selected from nucleotides 318-1766 of SEQ ID NO:15, at least 11 contiguous nucleotides selected from nucleotides 1-42 of SEQ ID NO:15, at least 11 contiguous nucleotides selected from nucleotides 478-908 of SEQ ID NO:15, at least 11 contiguous nucleotides selected from nucleotides 1059-1078 of SEQ ID NO:15, at least 205 contiguous nucleotides of SEQ ID NO:17, at least 440 contiguous nucleotides of SEQ ID NO:19, at least 451 contiguous nucleotides of SEQ ID NO:21, at least 11 contiguous nucleotides selected from nucleotides 1-121 of SEQ ID NO:21, at least 11 contiguous nucleotides selected from nucleotides 474-592 of SEQ ID NO:21, at least 351 contiguous nucleotides of SEQ ID NO:23, at least 21 contiguous nucleotides selected from nucleotides 1-1943 of SEQ ID NO:23, at least 11 contiguous nucleotides selected from 1-612 of SEQ ID NO:23, at least 11 contiguous nucleotides selected from nucleotides 611-719 of SEQ ID NO:23, at least 11 contiguous nucleotides selected from nucleotides 713-830 of SEQ ID NO:23, at least 11 contiguous nucleotides selected from nucleotides 830-1933 of SEQ ID NO:23, at least 492 nucleotides of SEQ ID NO:25, at least 11 contiguous nucleotides selected from nucleotides 758-847 of SEQ ID NO:25, at least 1024 contiguous nucleotides of SEQ ID NO:27, at least 347 contiguous nucleotides of SEQ ID NO: 29, at least 11 contiguous nucleotides selected from nucleotides 548-601 of SEQ ID NO:29, at least 394 contiguous nucleotides of SEQ ID NO: 31, at least 11 contiguous nucleotides selected from nucleotides 1-361 of SEQ ID NO:31, at least 11 contiguous nucleotides selected from nucleotides 1083-1102 of SEQ ID NO:31, at least 492 contiguous nucleotides of SEQ ID NO:33, at least 510 contiguous nucleotides of SEQ ID NO:35, at least 11 contiguous nucleotides selected from nucleotides 1-502 or 505-631 of SEQ ID NO:35, at least 392 contiguous nucleotides of SEQ ID NO:37, at least 11 contiguous nucleotides selected from nucleotides 1-502 of SEQ ID NO:37, at least 11 contiguous nucleotides selected from nucleotides 505-631 of SEQ ID NO:37, at least 559 contiguous nucleotides of SEQ ID NO:39, at least 11 contiguous nucleotides selected from nucleotides 1-92 of SEQ ID NO:39, at least 254 contiguous nucleotides of SEQ ID NO:41, at least 11 contiguous

nucleotides selected from nucleotides 1-34 of SEQ ID NO:41 at least 11 contiguous nucleotides selected from nucleotides 55-110 of SEQ ID NO:41, at least 103 contiguous nucleotides of SEQ ID NO:43, at least 11 contiguous nucleotides selected from nucleotides 1-280 of SEQ ID NO:43, at least 11 contiguous nucleotides selected from nucleotides 270-319 of SEQ ID NO:43, at least 11 contiguous nucleotides selected from nucleotides 378-423 of SEQ ID NO:43, at least 11 contiguous nucleotides selected from nucleotides 414-492 of SEQ ID NO:43, at least 11 contiguous nucleotides selected from nucleotides 532-570 of SEQ ID NO:43, at least 11 contiguous nucleotides selected from nucleotides 1086-1152 of SEQ ID NO:43, and the complements thereof.

14. A construct comprising the isolated and purified subgenomic polynucleotide of claim 9.
15. The construct of claim 14 further comprising a promoter which is operatively linked to the nucleotide sequence.
16. A host cell comprising the construct of claim 14.
17. The host cell of claim 16 which is a mammalian cell.
18. A process for producing a protein, comprising the steps of:
  - growing a culture of the host cell of claim 66 in a suitable culture medium; and
  - purifying the protein secreted from the host cell.
19. A polynucleotide array comprising at least one single-stranded polynucleotide which comprises at least 12 contiguous nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, and 43.

20. A method of detecting differential gene expression between two biological samples, comprising the step of:

contacting a first biological sample comprising single-stranded polynucleotide molecules with a first polynucleotide array comprising at least one single-stranded polynucleotide which comprises at least 12 contiguous nucleotides of a nucleotide sequence selected from the group consisting of SEQ ID NOS:1, 3, 5, 7, 9, 11, 13, 15, 17, 19, 21, 23, 25, 27, 29, 31, 33, 35, 37, 39, 41, and 43;

contacting a second biological sample comprising single-stranded polynucleotide molecules with a second polynucleotide array, wherein the first and second polynucleotide arrays comprise identical single-stranded polynucleotides; and

detecting a first and second pattern of double-stranded polynucleotides bound to the first and second polynucleotide arrays, wherein a difference between the first and second patterns indicates a gene which is differentially expressed between the first and second biological samples.

21. The method of claim 20 wherein the first biological sample is suspected of being diseased and wherein the second biological sample is not diseased.

**SEQUENCE LISTING**

SEQ ID NO:1 (hCornichon cDNA)

```

      10      20      30      40      50      60
GTTACAGTTCGCGGCCTTCTGCTACATGCTGGCGCTGCTGCTCACTGCCGCGCTCATCTT

      70      80      90     100     110     120
CTTCGCCATTTGGCACATTATAGCATTTGATGAGCTGAAGACTGATTACAAGAATCCTAT

     130     140     150     160     170     180
AGACCAGTGTAAATACCCCTGAATCCCCTTGTA TCTCCCAGAGTACCTCATCCACGCTTCTT

     190     200     210     220     230     240
CTGTGTCATGTTTCTTTGTGTCAGCAGAGTGGCTTACTGGGTCTCAATATGCCCTCTT

     250     260     270     280     290     300
GGCATATCATATTTGGAGGTATATGAGTAGACCAGTGATGAGTGGCCAGGACTCTATGA

     310     320     330     340     350     360
CCCTACAACCATCATGAATGCAGATATTCTAGCATATTGTCAGAAGGAAGGATGGTGCAA

     370     380     390     400     410     420
ATTAGCTTTTTATCTTCTAGCATTTTTTTACTACCTATATGGCATGATCTATGTTTGGT

     430     440     450     460     470     480
GAGCTCTTAGAACAACACACAGAAGAATTGGTCCAGTTAAGTGCATGCAAAAAGCCACCA

     490     500     510     520     530     540
AATGAAGGGATTCTATCCAGCAAGATCCTGTCCAAGAGTAGCCTGTGGAATCTGATCAGT

     550     560     570     580     590     600
TACTTTAAAAAATGACTCCTTATTTTTTAAATGTTCCACATTTTGTGTTGTGAAAGAC

     610     620     630     640     650     660
TGTTTTCATATGTTATACTCAGATAAAGATTTTAAATGGTATTACGTATAAATTAATATA

     670     680     690     700     710     720
AAATGGTTACCTCTGGTGTGACAGGTTTGAACCTTGCACCTCTTAAGGAACAGCCATAAT

     730     740     750     760     770     780
CCTCTGAATGATGCATTAATTACTGACTGTCCCTAGTACATTGGAAGCTTTTGTTTATAGG

     790     800     810     820     830     840
AACTTGTAGGGCTCATTTTGGTTTCATTGAAACAGTATCTAATTATAAATTAGCTGTAGA

     850     860     870     880     890     900
TATCAGGTGCTTCTGATGAAGTGAAATGTATATCTGACTAGTGGGAAACTTCATGGGTT

     910     920     930     940     950     960
TCCTCATCTGTGTCATGTCGATGATTATATATGGATACATTTACAAAAATAAAAAGCGGGAA

     970     980     990    1000    1010    1020
TTTTCCCTTCGCTTGAATATTATCCCTGTATATTGCATGAATGAGAGATTTCCCATATTT

    1030    1040    1050    1060    1070    1080
CCATCAGAGTAATAAATATACTTGCTTTAATTCTTAAGCATAAGTAAACATGATATAAAA

    1090    1100    1110    1120    1130    1140
ATATATGCTGAATTACTTGTGAAGAATGCATTAAAGCTATTTTAAATGTGTTTTATT

    1150    1160    1170    1180    1190    1200
GTAAGACATTACTTATTAAGAAATTGGTTATTATGCTTACTGTTCTAATCTGGTGGTAA

    1210    1220    1230    1240    1250    1260
GGTATTCTTAAGAATTGCAGGTACTACAGATTTTCAAACTGAATGAGAGAAAATTGTA

    1270    1280    1290    1300    1310    1320
TAACCATCTGCTGTTCTTTAGTGCAATACATAAACTCTGAAATTAAGACTCAAAAA

```

AAAAA

## SEQ ID NO:2 (hCornichon polypeptide)

```

      10      20      30      40      50      60
ETFAAFCYMLALLLTAALIEFFAIWHIIAFDELKTDYKNPIDQCNTLNPLVLPEYLIHAFF
      70      80      90     100     110     120
CVMFLCAAEWLTGLNMPLLAYHIWRYMSRPVMSGPGLYDPTTIMNADILAYCQKEGWCK
      130     140
LAFYLLAFFYYLYGMIYVLVSS

```

## SEQ ID NO:3 (BMS46 cDNA)

```

      10      20      30      40      50      60
CACGAGGAAACCAACGAGGGGACGCGGCCGAGGAGGTCGCTGCCACCGGGGGCGTGG
      70      80      90     100     110     120
GAGTGAGGTACCAGATTACAGCCATTGGCCCCGACGCCCTCTGTTCTCGGAATCCGGGTG
      130     140     150     160     170     180
CTGCGGATTGAGGTCCCGGTTCTAACGGTGGGATCGGTGTCCTCGGGATGAGATTGGC
      190     200     210     220     230     240
GTTTCCTCGGGGCTTTGGTGGGATCGGTGTCCTCAGGATGAGATTAGGGTTCTCTCGGG
      250     260     270     280     290     300
GCTTTCGGGATCTTACCTAATATCCGGTATTATTTATGAGAGGAGTGGTCTTGGCTGT
      310     320     330     340     350     360
CAGAACTGGATCCCTGGGGTGATATTGGGAATTAGTGGAGTGATCTCTGAAGACCTAGG
      370     380     390     400     410     420
GCTATGATCTGGAGCTGCTGTGGCTGAAATTTGGGGCCTCTGAAGTGGCATGGAGATTGA
      430     440     450     460     470     480
GGTCCAGAGAGCCTGAGATCTTGAGGGCTGACATTGGAGAGATGGGGTCGAGGGTTGTC
      490     500     510     520     530     540
TTTGGGCCTTGACTGCTTTGGGCCTTTCTCACTCTCATTCCCGGGATGCTTTGCCAGAAT
      550     560     570     580     590     600
CTCTGCTGGATTGGCCGTAAACCTGTCCCCGAGCGGGCTCACAGGGTCTGAAGGCCACGC
      610     620     630     640     650     660
ATGAGGCAAAGGTAAGTTCTGAGCCACCCGGTGCTCCTTCCAGGACTGCAAGATGGA
      670     680     690     700     710     720
GGAAGGCGGGAACCTAGGAGGCCTGATTAAGATGGTCCATCTACTGGTCTTGTCAGGTGC
      730     740     750     760     770     780
CTGGGGCATGCAAATGTGGGTGACCTCGTCTCAGGCTTCCTGCTTTCCGAAGCCTTCC
      790     800     810     820     830     840
CCGACATACCTTCGGACTAGTGCAGAGCAAACCTCTCCCTTCTACTTCCACATCTCCAT
      850     860     870     880     890     900
GGGCTGTGCCTTCATCAACCTCTGCATCTTGGCTTCACAGCATGCTTGGGCTCAGCTCAC
      910     920     930     940     950     960
ATTCTGGGAGGCCAGCCAGCTTTACCTGCTGTTCTGAGCCCTACGCTGGCCACTGTCAA
      970     980     990    1000    1010    1020
CGCCCGCTGGCTGGAACCCCGCACACAGCTGCCATGTGGGCCCTGCAAACCGTGGAGAA
      1030    1040    1050    1060    1070    1080
GGAGCGAGGCCTGGGTGGGGAGGTACCAGGCAGCCACCAGGGTCCCGATCCCTACCGCAA
      1090    1100    1110    1120    1130    1140
GCTGCGAGAGAAGGACCCCAAGTACAGTGCTCTCGCCAGAATTTCTTCGGCTACCATGG

```

1150 1160 1170 1180 1190 1200  
 GCTGTCCTCTCTTTGCAATCTGGGCTGCGTCCTGAGCAATGGGCTCTGTCTCGCTGGCCT  
 1210 1220 1230 1240 1250 1260  
 TGCCCTGGAAATAAGGAGCCTCTAGCATGGGCCCTGCATGCTAATAAATGCTTCTTCAGA  
 1270  
 AAAAAAAAAAAAAAAAAA

## SEQ ID NO:4 (BMS46 polypeptide)

10 20 30 40 50 60  
 MEEGGNIGGLIKMYHLLVLSGAWGMOMVTEVSGELLFRSLPRHTFGLVQSKLFFFYFHI  
 70 80 90 100 110 120  
 SMGCAFINLCLILASQHAWAQLTFWEASQLYLLFLSLTLATVNARWLEPRTTAAMWALQTV  
 130 140 150 160 170 180  
 EKERGLGGEVPGSHQGPDPYRQLREKDPKYSALRQNFYRYHGLSSLCNLGCVLSNGLCLA  
 GLALEIRSL

## SEQ ID NO:5 (BMS112 cDNA)

10 20 30 40 50 60  
 CACAGTAGGTCCCTCGGCTCAGTCGGCCCAGCCCCTCTCAGTCCTCCCAACCCCCACAA  
 70 80 90 100 110 120  
 CCGCCCGCGGCTCTGAGACGCGGCCCCGCGGCGGCGGCAGCAGCTGCAGCATCATCTCC  
 130 140 150 160 170 180  
 ACCCTCCAGCCATGGAAGACCTGGACCAGTCTCCTCTGGTCTCGTCCTCGGACAGCCCAC  
 190 200 210 220 230 240  
 CCCGGCCGCGAGCCCGGTTCAAGTACCAGTTCGTGAGGGAGCCCGAGGACGAGGAGGAAG  
 250 260 270 280 290 300  
 AAGAGGAGGAGGAAGAGGAGGACGAGGACGAAGACCTGGAGGAGCTGGAGGTGCTGGAGA  
 310 320 330 340 350 360  
 GGAAGCCCGCCCGGGCTGTCCGCGGCCCCAGTGCCACCGCCCCTGCCGCCGGCGCGC  
 370 380 390 400 410 420  
 CCCTGATGGACTTCGGAATGACTTCGTGCCGCGGCCCGGGGACCCCTGCCGGCCG  
 430 440 450 460 470 480  
 CTCCCCCGTCGCCCCGGAGCGGCAGCCGTCTTGGGACCGAGCCCGGTGTCGTCGACCG  
 490 500 510 520 530 540



TGCCCCGCGCCATCCCCGCTGTCTGCTGCGCAGTCTCGCCCTCCAAGCTCCCTGAGGACG  
550 560 570 580 590 600  
ACGAGCCTCCGGCCCCGGCCTCCCCCTCCTCCCCGGCCAGCGTGAGCCCCCAGGCAGAGC  
610 620 630 640 650 660  
CCGTGTGGACCCCGCCAGCCCCGGCTCCGCGCGCCCCCTCCACCCCGGCGCGCCCA  
670 680 690 700 710 720  
AGCGCAGGGGCTCCTCGGGCTCAGTGGTTGTTGACCTCCTGTACTGGAGAGACATTAAGA  
730 740 750 760 770 780  
AGACTGGAGTGGTGTGTTGGTGCCAGCCTATTCTGCTGCTTTCATTGACAGTATTCAGCA  
790 800 810 820 830 840  
TTGTGAGCGTAACAGCCTACATTGCCTTGCGCCCTGCTCTCTGTGACCATCAGCTTTAGGA  
850 860 870 880 890 900  
TATACAAGGGTGTGATCCAAGCTATCCAGAAATCAGATGAAGGCCACCCATTACAGGGCAT  
910 920 930 940 950 960  
ATCTGGAATCTGAAGTTGCTATATCTGAGGAGTTGGTTCAGAAGTACAGTAATTCTGCTC  
970 980 990 1000 1010 1020  
TTGGTCATGTCAACTGCACGATAAAGGAACTCAGGCGCCTCTTCTTAGTTGATGATTTAG  
1030 1040 1050 1060 1070 1080  
TTGATTCTCTGAAGTTTGCAGTGTGATGTGGGTATTTACCTATGTTGGTGCCTTGTTTA  
1090 1100 1110 1120 1130 1140  
ATGGTCTGACACTACTGATTTTGGCTCTCATTTCACTCTTCAGTGTTCTGTATTATG  
1150 1160 1170 1180 1190 1200  
AACGGCATCAGGCACAGATAGATCATTATCTAGGACTTGCAAATAAGAATGTTAAAGATG  
1210 1220 1230 1240 1250 1260  
CTATGGCTAAATCCAAGCAAAATCCCTGGATTGAAGCGCAAAGCTGAATGAAACGCC  
1270 1280 1290 1300 1310 1320  
CAAAATAATTAGTAGGAGTTCATCTTTAAAGGGGATATTCATTTGATTATACGGGGGAGG

1330 1340 1350 1360 1370 1380  
GTCAGGGAAGAACGAACCTTGACGTTGCAGTGCAGTTTCACAGATCGTTGTTAGATCTTT  
1390 1400 1410 1420 1430 1440  
ATTTTGTAGCCATGCACTGTTGTGAGGAAAAATTACCTGTCTTGACTGCCATGTGTTTCATC  
1450 1460 1470 1480 1490 1500  
ATCTTAAGTATTGTAAGCTGCTATGTATGGATTTAAACCGTAATCATATCTTTTTCCTAT  
1510 1520 1530 1540 1550 1560  
CTGAGGCACTGGTGGAAATAAAAAACCTGTATATTTTACTTTGTTGCAGATAGTCTTGCCG  
1570 1580 1590 1600 1610  
CATCTTGGCAAGTTGCAGAGATGGTGGAGCTAGAAAAAAAAAAAAAAAAAAAA

## SEQ ID NO:6 (BMS112 polypeptide)

10 20 30 40 50 60  
MEDLDQSPLVSSSDSPPRPQPAFKYQFVREPEDEEEEEEEEEDEDEDLEELEVLERKPA  
70 80 90 100 110 120  
AGLSAAPVPTAPAAGAPLMDFGND FVPPAPRGPLPAAPPVAPERQPSWDPSPVSSTVPAP  
130 140 150 160 170 180  
SPLSAAAVSPSKLPEDDEPPARPPPPPPASVSPQAEFVWTPPAPAPAAPPSTPAAPKRRG  
190 200 210 220 230 240  
SSGSVVVDLLYWRDIKKTGVVFGASLFLLLSLTVFSIVSVTAYIALALLSVTISFRIYKG  
250 260 270 280 290 300  
VIQAIQKSDEGHPFRAYLESEVAISEELVQKYSNSALGHVNCTIKELRRLFLVDDLVDLSL  
310 320 330 340 350 360  
KFAVLMWVFTYVGALFNGLTLLILALISLFSVPVIYERHQAQIDHYLGLANKNVKDAMAK  
370  
IQAKIPGLKRKAE

## SEQ ID NO:7 (BMS118 cDNA)

10 20 30 40 50 60  
GTCGAGAGGACGAGGTGCCGCTGCCTGGAGAATCCTCCGCTGCCGTCGGCTCCCGGAGCC

70 80 90 100 110 120  
CAGCCCTTTCTTAACCCAACCAACCTAGCCCAGTCCCAGCCGCCAGCGCCTGTCCCTGT

130 140 150 160 170 180  
CACGGACCCAGCGTTACCATGCATCCTGCCGTCTTCCTATCCTTACCCGACCTCAGATG

190 200 210 220 230 240  
CTCCCTTCTGCTCCTGGTAACTTGGGTTTTTACTCCTGTAACAACTGAAATAACAAGTCT

250 260 270 280 290 300  
TGATACAGAGAATATAGATGAAATTTTAAACAATGCTGATGTTGCTTTAGTAAATTTT

310 320 330 340 350 360  
TGCTGACTGGTGTCGTTTCAGTCAGATGTTGCATCCAATTTTGTAGGAAGCTTCCGATGT

370 380 390 400 410 420  
CATTAAGGAAGAATTTCCAAATGAAAATCAAGTAGTGTGTTGCCAGAGTTGATTGTGATCA

430 440 450 460 470 480  
GCACTCTGACATAGCCCAGAGATACAGGATAAGCAAATACCCAACCCTCAAATTGTTTCG

490 500 510 520 530 540  
TAATGGGATGATGATGAAGAGAGAATACAGGGGTCAGCGATCAGTGAAGCATTGGCAGA

550 560 570 580 590 600  
TTACATCAGGCAACAAAAAGTGACCCCATTCAGAAATTCGGGACTTAGCAGAAATC

610 620 630 640 650 660  
CACTCTTGATCGCAGCAAAAGAAATATCATTGGATATTTGAGCAAAAGGACTCGGACAA

670 680 690 700 710 720  
CTATAGAGTTTTTGAACGAGTAGCGAATATTTGCATGATGACTGTGCCTTTCTTTCTGC

730 740 750 760 770 780  
ATTTGGGGATGTTTCAAAACCGGAAGATATAGTGGCGACAACATAATCTACAAACCACC

790 800 810 820 830 840  
AGGGCATTCTGCTCCGGATATGGTGTACTTGGGAGCTATGACAAATTTTGATGTGACTTA

850 860 870 880 890 900  
CAATTGGATTCAAGATAAATGTGTTCCTCTGTCCGAGAAATAACATTTGAAAATGGAGA

910 920 930 940 950 960  
GGAATTGACAGAAGAAGGACTGCCTTTTCTCATACTCTTTCACATGAAAGAAGATACAGA

970 980 990 1000 1010 1020  
AAGTTTAGAAATATTCCAGAATGAAGTAGCTCGGCAATTAATAAGTGAAAAAGGTACAAT

1030 1040 1050 1060 1070 1080  
AAACTTTTTACATGCCGATTGTGACAAATTTAGACATCCTCTTCTGCACATACAGAAAAC

1090 1100 1110 1120 1130 1140  
TCCAGCAGATTGTCCTGTAATCGCTATTGACAGCTTTAGGCATATGTATGTGTTGGAGA

1150 1160 1170 1180 1190 1200  
CTTCAAAGATGTATTAATTCCTGGAAACTCAAGCAATTCGTATTTGACTTACATTCTGG

1210 1220 1230 1240 1250 1260  
AAAAGTGCACAGAGAATTCCATCATGGACCTGACCCAACTGATACAGCCCCAGGAGAGCA

1270 1280 1290 1300 1310 1320  
AGCCCCAAGATGTAGCAAGCAGTCCACCTGAGAGCTCCTTCCAGAACTAGCACCCAGTGA

1330 1340 1350 1360 1370 1380  
ATATAGGTATACTCTATTGAGGGATCGAGATGAGCTTTAAAAACTTGAAAAACAGTTTGT

1390 1400 1410 1420 1430 1440  
AAGCCTTTCAACAGCAGCATCAACCTACGTGGTGGAATAGTAAACCTATATTTTCATAA

1450 1460 1470 1480 1490  
TTCTATGTGTATTTTTATTTTGAATAAACAGAAAGAAATTTAAAAAAAAAAAAAAAAAAAA

## SEQ ID NO:8 (BMS118 polypeptide)

10	20	30	40	50	60
MHPAVFLSLPDLRCSLLLLVTWVTPVTEITSLDTENIDEILNNADVALVNFYADWCRF					
70	80	90	100	110	120
SQMLHPIFEEASDVIKEEFPNENQVVFARVDCDQHS DIAQRYRISKYPTLKLFRNGMMK					
130	140	150	160	170	180
REYRGQRSVKALADYIRQKSDPIQEIRDLAEITTLDRSKRNIIGYFEQKDSDNRYRVER					
190	200	210	220	230	240
VANILHDDCAFLSAFGDVSKPERYSGDNIIYKPPGHSAPDMVYLGAMTNFDVTYNWIQDK					
250	260	270	280	290	300
CVPLVREITFENGEEELTEEGLPFLILFHKEDTESLEIFQNEVARQLISEKGTINFLHAD					
310	320	330	340	350	360
CDKFRHPLLHIQKTPADCPVIAIDSFRHMYVFGDFKDVLI PGKLKQFVFDLHSGKLHREF					
370	380	390	400		
HHGPDPTDTAPGEQAQDVASSPPESSFQKLAPSEYRYTLLRDRDEL					

## SEQ ID NO:9 (BMS164 cDNA)

10	20	30	40	50	60
GCCTTTCCGCGCTTCTGCCGTGGCCCTCTGCCGGCCGCTCCGCCGGTGCTGTCCCTGGGCG					
70	80	90	100	110	120
CCTCCGTGCTCTCAGCCAACCGCCTCTGAGAGCGCCCACTCGAGCGCCCCGGGAGCCAGA					
130	140	150	160	170	180
GGGCGGGGGTCTCGCCGGGACCTCCTGTGGGCCCAGGGGGACAAAAGTGGCTCTCAAT					
190	200	210	220	230	240
CCAGCACATGCACATTGAAGCAAGTTAAAGGATTTAATATGAAGCACAGAAGCAGATAGT					
250	260	270	280	290	300
GCCAAATAGCAAGCAGTAGTTGTTACACATTTGGTGAGCAGGGCAGCATTTCCTTCTCCC					

310 320 330 340 350 360  
ACTGCTGCTGAGATGGCAGAAATTAGTCGAATTCAGTACGAAATGGAATATACTGAAGGC

370 380 390 400 410 420  
ATTAGTCAGCGAATGAGGGTCCCAGAAAAGTTAAAAGTAGCACCGCCAAACGCTGACCTG

430 440 450 460 470 480  
GAACAAGGATTCCAAGAAGGAGTTCCAAATGCTAGTGTGATAATGCAAGTTCCGGAGAGG

490 500 510 520 530 540  
ATTGTTGTAGCAGGAAATAATGAAGATGTTTCATTTTCAAGACCAGCAGATCTTGACCTT

550 560 570 580 590 600  
ATTCAGTCAACTCCCTTTAAACCCCTGGCACTGAAAACACCACCTCGTGTACTTACGCTG

610 620 630 640 650 660  
AGTGAAAGACCACTAGATTTTCTGGATTTAGAAAGACCTCCTACAACCCCTCAAATGAA

670 680 690 700 710 720  
GAAATCCGAGCAGTTGGCAGACTAAAAAGAGAGCGGTCTATGAGTGAAAATGCTGTTTCGC

730 740 750 760 770 780  
CAAATGGACAGCTGGTCAGAAATGATTCTTCTGTGACACCATCGCCACAACAGGCTCGG

790 800 810 820 830 840  
GTCTGTCTCTCCCATATGTTACCTGAAGATGGAGCTAATCTTCTCTGCTCGTGGCATT

850 860 870 880 890 900  
TTGTGCTTATCCAGTCTTCTACTCGTAGGGCATAACCAGCAGATCTTGGATGTGCTGGAT

910 920 930 940 950 960  
GAAAATCGCAGACCTGTGTTGCGTGGTGGGTCTGCTGCCGCCACTTCTAATCCTCATCAT

970 980 990 1000 1010 1020  
GACAACGTCAGGTATGGCATTTCAAATATAGATACAACCATTGAAGGAACGTCAGATGAC

1030 1040 1050 1060 1070 1080  
CTGACTGTTGTAGATGCAGCTTCACTAAGACGACAGATAATCAAACCTAAATAGACGTCTA

1090      1100      1110      1120      1130      1140  
 CAACTTCTGGAAGAGGAGAACAAGAACGTGCTAAAAGAGAAATGGTCATGTATTCAATT  
 1150      1160      1170      1180      1190      1200  
 ACTGTAGCTTTCTGGCTGCTTAATAGCTGGCTCTGGTTTCGCCGCTAGAGGTAACATCAG  
 1210      1220      1230      1240      1250      1260  
 CCCTCAAAAATACTGTCTCAACAGCTGGAAATATAAAAGATTGCAAACCTCAAAAAAAA  
 1270  
 AAAAAAAAAA

SEQ ID NO:10 (BMS164 polypeptide)

10      20      30      40      50      60  
 MAEISRIQYEMEYTEGISQRMRVPEKLKVAPPNADLEQGFQEGVPNASVIMQVPERIVVA  
 70      80      90      100      110      120  
 GNNEDVSFSRPADLDLIQSTPFKPLALKTPPRVLTLSERPLDFDLERPPTTPQNEEIRA  
 130      140      150      160      170      180  
 VGR LKRERSMSENAVRQNGQLVRNDSLVTSPQQARVCPPHMLPEDGANLSSARGILSLI  
 190      200      210      220      230      240  
 QSSTRRAYQQILDVLDENRRPVLRGGSAAATSNPHHDNVRYGISNIDTTIEGTSDDLTVV  
 250      260      270      280      290  
 DAASLRQIIKLNRRQLLEEENKERAKREVMYSITVAFWLLNSWLWFR

SEQ ID NO:11 (BMS192 cDNA)

10      20      30      40      50      60  
 GCGGCCCGGGCGGGCTGCTCGGCGCGGAACAGTGCTCGGCATGGCAGGGATTCCAGGGCT  
 70      80      90      100      110      120  
 CCTCTTCCTTCTCTTCTGCTCTGTGCTGTTGGGCAAGTGAGCCCTTACAGTGCCCC  
 130      140      150      160      170      180  
 CTGGAAACCCACTTGGCCTGCATACCGCCTCCCTGTCGTCTTGCCCCAGTCTACCCCTCAA  
 190      200      210      220      230      240

TTTAGCCAAGCCAGACTTTGGAGCCGAAGCCAAATTAGAAGTATCTTCTTCATGTGGACC  
 250 260 270 280 290 300  
 CCAGTGTCTATAAGGGAAGTCCACTGCCCCTTACGAAGAGGCCAAGCAATATCTGTCTTA  
 310 320 330 340 350 360  
 TGAAACGCTCTATGCCAATGGCAGCCGCACAGAGACGCAGGTGGGCATCTACATCCTCAG  
 370 380 390 400 410 420  
 CAGTAGTGGAGATGGGGCCCAACACCGAGACTCAGGGTCTTCAGGAAAGTCTCGAAGGAA  
 430 440 450 460 470 480  
 GCGGCAGATTTATGGCTATGACAGCAGGTTTCAGCATTTTGGGAAGGACTTCCTGCTCAA  
 490 500 510 520 530 540  
 CTACCCTTTCTCAACATCAGTGAAGTTATCCACGGGCTGCACCGGCACCCTGGTGGCAGA  
 550 560 570 580 590 600  
 GAAGCATGTCCTCACAGCTGCCCCTGCATACACGATGGAAAACCTATGTGAAAGGAAC  
 610 620 630 640 650 660  
 CCAGAAGCTTCGAGTGGGCTTCCTAAAGCCCAAGTTTAAAGATGGTGGTTCGAGGGGCCAA  
 670 680 690 700 710 720  
 CGACTCCACTTCAGCCATGCCCGAGCAGATGAAATTTTCAGTGGATCCGGGTGAAACGCAC  
 730 740 750 760 770 780  
 CCATGTGCCCAGGGTTGGATCAAGGGCAATGCCAATGACATCGGCATGGATTATGATTA  
 790 800 810 820 830 840  
 TGCCCTCCTGGAAGTCAAAAAGCCCCACAAGAGAAAATTTATGAAGATTGGGGTGAGCCC  
 850 860 870 880 890 900  
 TCCTGCTAAGCAGCTGCCAGGGGGCAGAATTCACCTTCTCTGGTTATGACAATGACCGACC  
 910 920 930 940 950 960  
 AGGCAATTTGGTGTATCGCTTCTGTGACGTCAAAGACGAGACCTATGACTTGCTCTACCA  
 970 980 990 1000 1010 1020  
 GCAATGCGATGCCAGCCAGGGGCCAGCGGGTCTGGGGTCTATGTGAGGATGTGGAAGAG



SEQ ID NO: 12 (BMS192 polypeptide)

12

130 140 150 160 170 180  
SGKSRRKRQIYGYSRFSIFGKDFLLNYPFSTSVKLSTGCTGTLVAEKHVLTAACIHDG  
190 200 210 220 230 240  
KTYVKGTTQKL RVGFLKPKFKDGGRGANDSTSAMPEQMKFQWIRVKRTHVPKGWIKGNAND  
250 260 270 280 290 300  
IGMDYDYALLELKKPHKRKFMKIGVSPPAKQLPGGRIHFSGYDNDRPGNLVYRFCDVKDE  
310 320 330 340 350 360  
TYDLLYQQCDAQPGASGSGVYVRMWKRQQQKWERKIIGIFSGHQWVDMNGSPQDFNVAVR  
370 380  
ITPLKYAQICYWIKGNYLDCREG

## SEQ ID NO:13 (BMS227 cDNA)

10 20 30 40 50 60  
CAGTAAGCTCGGCTCACAGTCGCAGGAGAGTTCTGGGGTACACGGGCAAAGGGGCTTGAG  
70 80 90 100 110 120  
AAGGCCCCGAGGCGAAGCCGAAGAGAAGCAACTGTGCCCCGAGAGAAGAGCTCGCCCA  
130 140 150 160 170 180  
TTCCAGACTGGGAACCAGCTTTCAGTGAAGATGGCAGGGCCAGAACTGTTGCTCGACTCC  
190 200 210 220 230 240  
AACATCCGCCTCTGGGTGGTCCTACCCATCGTTATCATCACTTCTTCGTAGGCATGATC  
250 260 270 280 290 300  
CGCCACTACGTGTCCATCCTGCTGCAGAGCGACAAGAAGCTCAGGAGGAAACAAGTATCT  
310 320 330 340 350 360  
GACAGTCAAGTCCTAATTCGAAGCAGAGTCCTCAGGGAAAATGGAAAATACATTCCCAAA  
370 380 390 400 410 420  
CAGTCTTTCTTGACACGAAAATATTATTTCAACAACCCAGAGGATGGATTTTTCAAAAA  
430 440 450 460 470 480

ACTAAACGGAAGGTAGTGCCACCTTCTCCTATGACTGATCCTACTATGTTGACAGACATG  
490 500 510 520 530 540  
ATGAAAGGGAATGTAACAAATGTCCTCCCTATGATTCTTATTGGTGGATGGATCAACATG  
550 560 570 580 590 600  
ACATTCTCAGGCTTTGTGACAACCAAGGTCCCATTTCCACTGACCCTCCGTTTAAAGCCT  
610 620 630 640 650 660  
ATGTTACAGCAAGGAATCGAGCTACTCACATTAGATGCATCCTGGGTGAGTTCTGCATCC  
670 680 690 700 710 720  
TGGTACTTCCTCAATGTATTTGGGCTTCGGAGCATTACTCTCTGATTCTGGGCCAAGAT  
730 740 750 760 770 780  
AATGCCGCTGACCAATCACGAATGATGCAGGAGCAGATGACGGGAGCAGCCATGGCCATG  
790 800 810 820 830 840  
CCCGCAGACACAAACAAAGCTTTCAAGACAGAGTGGGAAGCTTTGGAGCTGACGGATCAC  
850 860 870 880 890 900  
CAGTGGGCACTAGATGATGTGGAAGAAGAGCTCATGGCCAAAGACCTCCACTTCGAAGGC  
910 920 930 940 950 960  
ATGTTCAAAAAGGAATTACAGACCTCTATTTTTTGAAGACCGAGCAGGGATTAGCTGTGT  
970 980 990 1000 1010 1020  
CAGGAAGTTGGAGTTGCACTTAACCTTGTAACCTTTGTTTGGAGCTGGCACCTCTTGAAT  
1030 1040 1050 1060 1070  
AAAAAGGAGGATGCACGAGCTGGCAGGCATGCAAAAAAAAAAAAAAAAAAAAA

SEQ ID NO:14 (BMS227 polypeptide)

10 20 30 40 50 60  
MAGPELLLDENIRLWVVLPIVLIITFFVGMIRHYVSILLQSDKKLTQEQVSDSQVLIRSV  
70 80 90 100 110 120  
LRENGKYIPKQSFLTRKYYFNNPEDGFFKTKRKVVPPSPMTDPTMLTDMMKGNVTNVL

130 140 150 160 170 180  
MILIGGWINMTFSGFVTTKVPFPLTLRFKPHLQQGIELLTLDSWVSSASWYFLNVFGLR  
190 200 210 220 230 240  
SIYSLILGQDNAADQSRMMQEQMTGAAMAMPADTNKAFKTEWEALELTDHQWALDDVEEE  
250 260  
LMAKDLHFEGMFKKELQTSIF

SEQ ID NO:15 (BMS115 cDNA)

10 20 30 40 50 60  
ATGGCGGCCGCGGGGCTGCGGCTACACACCTAGAGGTGGCCCCGGGGCAAGCGCGCCGCC  
70 80 90 100 110 120  
CTCTTCTTCGCTGCGGTGGCCATCGTGCTGGGGCTACCGCTCTGGTGAAGACCACGGAG  
130 140 150 160 170 180  
ACCTACCGGGCCTCGTTGCCTTACTCCCAGATCAGTGGCCTGAATGCCCTTCAGCTCCGC  
190 200 210 220 230 240  
CTCATGGTGCCTGTCACTGTCGTGTTTACGCGGGAGTCAGTGCCCTGGACGACCAGGAG  
250 260 270 280 290 300  
AAGCTGCCCTTCACCGTTGTGCATGAAAGAGAGATTCTCTGAAATACAAAATGAAAATC  
310 320 330 340 350 360  
AAATGCCGTTTCCAGAAGGCCTATCGGAGGGCTTTGGACCATGAGGAGGAGGCCCTGTCA  
370 380 390 400 410 420  
TCGGGCAGTGTGCAAGAGGCAGAAGCCATGTTAGATGAGCCTCAGGAACAAGCGGAGGGC  
430 440 450 460 470 480  
TCCCTGACTGTGTACGTGATATCTGAACACTCCTCACTTCTCCCCAGGACATGATGAGC  
490 500 510 520 530 540  
TACATTGGGCCCCAAGAGGACAGCAGTGGTGGGGGGATAATGCACCGGGAGGCCTTTAAC

550 560 570 580 590 600  
ATCATTGGCCGCGCATAGTCCAGGTGGCCCAGGCCATGTCTTTGACTGAGGATGTGCTT

610 620 630 640 650 660  
GCTGCTGCTCTGGCTGACCACCTTCCAGAGGACAAGTGGAGCGCTGAGAAGAGGCGGCCT

670 680 690 700 710 720  
CTCAAGTCCAGCTTGGGCTATGAGATCACCTTCAGTTTACTCAACCCAGACCCCAAGTCC

730 740 750 760 770 780  
CATGATGTCTACTGGGACATTGAGGGGGCTGTCCGGCGCTATGTGCAACCTTTCCTGAAT

790 800 810 820 830 840  
GCCCTCGGTGCCGCTGGCAACTTCTCTGTGGACTCTCAGATTCTTTACTATGCAATGTTG

850 860 870 880 890 900  
GGGGTGAATCCCGCTTTGACTCAGCTTCCTCCAGCTACTATTTGGACATGCACAGCCTC

910 920 930 940 950 960  
CCCCATGTCAACCCAGTGGAGTCCCGGCTGGGATCCAGTGCTGCCTCCTTGTACCTT

970 980 990 1000 1010 1020  
GTGCTCAACTTTCTACTCTACGTGCCTGAGCTTGCACACTCACCGCTGTACATTCAGGAC

1030 1040 1050 1060 1070 1080  
AAGGATGGCGCTCCAGTGGCCACCAATGCCTTCCATAGTCCCGCTGGGGTGGCATTATG

1090 1100 1110 1120 1130 1140  
GTATATAATGTTGACTCCAAAACCTATAATGCCTCAGTGCTGCCAGTGAGAGTCGAGGTG

1150 1160 1170 1180 1190 1200  
GACATGGTGCGAGTGATGGAGGTGTTCTGGCACAGTTGCGGTTGCTCTTTGGGATTGCT

1210 1220 1230 1240 1250 1260  
CAGCCCCAGCTGCCTCCAAAATGCCTGCTTTTCAAGGCTACGAGTGAAGGGCTAATGACC

1270 1280 1290 1300 1310 1320  
TGGGAGCTAGACCGGCTGCTCTGGGCTCGGTGAGTGGAGAACCTGGCCACAGCCACCACC

1330 1340 1350 1360 1370 1380  
ACCCTTACCTCCCTGGCGCAGCTTCTGGGCAAGATCAGCAACATTGTCATTAAGGACGAC

1390 1400 1410 1420 1430 1440  
GTGGCATCTGAGGTGTACAAGGCTGTAGCTGCCGTCCAGAAGTCGGCAGAAGAGTTGGCG

1450 1460 1470 1480 1490 1500  
TCTGGGCACCTGGCATCTGCCTTTGTGCGCCAGCCAGGAAGCTGTGACATCCTCTGAGCTT

1510 1520 1530 1540 1550 1560  
GCCTTCTTTGACCCGTCACCTCCTCCACCTCCTTTATTTCCCTGATGACCAGAAGTTTGCC

1570 1580 1590 1600 1610 1620  
ATCTACATCCCACCTTCTCCTGCCTATGGCTGTGCCCATCCTCCTGTCCCTGGTCAAGATC

1630 1640 1650 1660 1670 1680  
TTCCTGGAGACCCGCAAGTCCTGGAGAAAGCCTGAGAAGACAGACTGAGCAGGGCAGCAC

1690 1700 1710 1720 1730 1740  
CTCCATAGGAAGCCTTCCTTTCTGGCCAAGGTGGGCGGTGTTAGATTGTGAGGCACGTAC

1750 1760 1770 1780 1790 1800  
ATGGGGCCTGCCGGAATGACTTAAATATTTGTCTCCAGTCTCCACTGTTGGCTCTCCAGC

1810 1820 1830 1840 1850 1860  
AACCRAAGTACAACACTCCAAGATGGGTTTCATCTTTTCTCCTTTCCCATTCACCTGGCT

1870 1880 1890 1900 1910 1920  
CAATCCTCCTCCACCACCAGGGGCCTCAAAGGCACATCATCCGGGTCTCCTTATCTTGT

1930 1940 1950 1960 1970 1980  
TTGATAAGGCTGCTGCCTGTCTCCCTCTGTGGCAAGGACTGTTTGTCTTTTGCCCCATT

1990 2000 2010 2020 2030 2040  
TCTCAACATAGCACACTTGTGCACTGAGAGGAGGGAGCATTATGGGAAAGTCCCTGCCTT

2050 2060 2070 2080 2090 2100  
CCACACCTCTCTCTAGTCCCTGTGGGACAGCCCTAGCCCCTGCTGTCATGAAGGGGCCAG

**SEQ ID NO:16 (BMS115 protein)**

18

310 320 330 340 350 360  
PHVINPVESRLGSSAASLYPVLNFLLYVPELAHSPLYIQDKDGAPVATNAFHSRWGGIM  
370 380 390 400 410 420  
VYNVDSKTYNASVLPVRVEVDMVRVMEVFLAQLRLLFGIAQPQLPPKCLLSGPTSEGLMT  
430 440 450 460 470 480  
WELDRLLWARSVENLATATTTLSLAQLLGKISNIVIKDDVASEVYKAAVQKSAEELA  
490 500 510 520 530 540  
SGHLASAFVASQEAVTSSELAFDPSSLHLLYFPDDQKFAIYIPLFLPMAVPILLSLVKI  
550  
FLETRKSWRKPEKTD

## SEQ ID NO:17 (BMS143 cDNA)

10 20 30 40 50 60  
CTACATCCTGGACAACGAGACCAACTTCGTGGTCCAGGTCAGCGTCTTCATTGGGGTCCT  
70 80 90 100 110 120  
CATCGACCTCTGGAAGATCACCAAGGTCATGGACGTCCGGCTGGACCGAGAGCACAGGGT  
130 140 150 160 170 180  
GGCAGGAATCTTCCCCGCCTATCCTTCAAGGACAAGTCCACGTATATCGAGTCCTCGAC  
190 200 210 220 230 240  
CAAAGTGTATGATGATATGGCATTCCGGTACCTGTCTGGATCCTCTCCCGCTCCTGGG  
250 260 270 280 290 300  
CTGCTATGCCGTCTACAGTCTTCTGTACCTGGAGCACAAAGGGCTGGTACTCCTGGGTGCT  
310 320 330 340 350 360  
CAGCATGCTCTACGGCTTCCTGCTGACCTTCGGCTTCATCACCATGACGCCCCAGCTCTT  
370 380 390 400 410 420  
CATCAACTACAAGCTCAAGTCTGTGGCCACCTTCCTGGCGCATGCTCACCTACAAGGC



430 440 450 460 470 480  
CCTCAACACATTTCATCGACGACCTGTTTCGCCTTTGTCATCAAGATGCCCCGTTATGTACCG

490 500 510 520 530 540  
GATCGGCTGCCTGCGGGACGATGTGGTTTTCTTCATCTACCTCTACCAACGGTGGATCTA

550 560 570 580 590 600  
CCGCGTCGACCCACCCGAGTCAACGAGTTTGGCATGAGTGGAGAAGACCCACAGCTGC

610 620 630 640 650 660  
CGCCCCCGTGGCCGAGGTTCCACAGCAGCAGGGGGCCCTCAGCCCCACACCTGCACCCAC

670 680 690 700 710 720  
CACGACCACCGCCACCAGGGAGGAGGCCTCCACGTCCCTGCCCACCAAGCCCCACCCAGGG

730 740 750 760 770 780  
GGCCAGCTCTGCCAGCGAGCCCCAGGAAGCCCCTCCAAAGCCAGCAGAGGACAAGAAAAA

790 800 810 820 830 840  
GGATTAGTCGAGACTGGTCCTCACCTGCTCCGGCTCCTGGCGACCACTACCCCTGCGTCC

850 860 870 880 890 900  
CGGCCCCCTCGCCTCCCCTCCCTGTGCGCCCTTCCCTGGACAGATCAGGCCGGGGCGGTG

910 920 930 940 950 960  
GGAGGCCCGCCTCAGGTGAGGGCCAGCGTGTGATGTAGGGGCCGGGGCAGGCCAGGGTT

970 980 990 1000 1010 1020  
TGTTTGTGGAGGCGCTGTCTGTCCCTCTGTCCCTCTGTGTTTCCAGCCATCTCGCCCTGC

1030 1040 1050 1060 1070 1080  
CAGCCCAGCACCCTGGGAATCATGGTGAAGCTGATGCAGCGTTGCCGAGGGGGTGGGTT

1090 1100 1110 1120 1130 1140  
GGGCGGGGGTGGGGCCGGGCCCCCTAGGGGATGCCCCGGGGCCGTTTCATCATCTTGTCCTC

1150 1160 1170 1180 1190 1200  
TGGTCCCCCTACCACACTCCCCCTCCTAAACCGCCGCCCTTTAACACAGTTTGGATTAA

1210 1220 1230 1240  
TAAATTTCAGATGGGGGTTTAACTTAACTCAAAAAAAAAAAAAA

## SEQ ID NO:18 (BMS143 protein)

10 20 30 40 50 60  
MDVRLDREHRVAGIEPRLSEFKDKSTYIESSTKVYDDMAFRYLSWILFPLLGCXAVYSLLY  
70 80 90 100 110 120  
LEHKGWYSWVLSMLYGFLLTFGFITMTPLQFIFYKLSVAHLPWRMLTYKALNTFIDDLF  
130 140 150 160 170 180  
AFVIKMPVMYRIGCLRDDVVFIFYLYQRWIYRVDPTRVNEFGMSGEDPTAAAPVAEVPTA  
190 200 210 220 230  
AGALTPTPAPTPTTTATREEASTSLPTKPTQGASSASEPQEAPPKPAEDKKKD

## SEQ ID NO:19 (BMS155 cDNA)

10 20 30 40 50 60  
AACATGGAGACTTTGTACCGTGTCCCGTTCTTAGTGCTCGAATGTCCCAACCTGAAGCTG  
70 80 90 100 110 120  
AAGAAGCCGCCCTGGTTCACATGCCGTCGGCCATGACTGTGTATGCTCTGGTGGTGGTG  
130 140 150 160 170 180  
TCTTACTTCCTCATCACCAGGAATAATTTATGATGTTATTGTTGAACCTCCAAGTGTC  
190 200 210 220 230 240  
GGTTCTATGACTGATGAACATGGGCATCAGAGGCCAGTAGCTTTCTTGGCCTACAGAGTA  
250 260 270 280 290 300  
AATGGACAATATATTATGGAAGGACTTGCATCCAGCTTCCTATTTACAATGGGAGGTTTA  
310 320 330 340 350 360  
GGTTTCATAATCCTGGACCGATCGAATGCACCAAATATCCCAAACTCAATAGATTCCCTT  
370 380 390 400 410 420  
CTTCTGTTTCATTGGATTCTGTGTCTTATTGAGTTTTTTCATGGCTAGAGTATTCATG  
430 440 450 460 470 480  
AGAATGAAACTGCCGGGCTATCTGATGGGTTAGAGTGCCTTTGAGAAGAAATCAGTGGAT

490 500 510 520 530 540  
ACTGGATTGCTCCTGTCAATGAAGTTTTAAAGGCTGTACCAATCCTCTAATATGAAATG  
550 560 570 580 590 600  
TGGAAAAGAATGAAGAGCAGCAGTAAAGAAATATCTAGTGAAAAACAGGAAGCGTATT  
610 620 630 640 650 660  
GAAGCTTGGACTAGAATTTCTTCTTGGTATTAAAGAGACAAGTTTATCACAGAATTTTTT  
670 680 690 700 710 720  
TTCCTGCTGGCCTATTGCTATACCAATGATGTTGAGTGGCATTTCCTTTTAGTTTTTCA  
730 740 750 760 770 780  
TTAAAATATATTCCATATCTACAACATAATATCAAATAAAGTGATTATTTTTTACAACC  
790 800 810 820 830 840  
CTCTTAACATTTTTTGGAGATGACATTTCTGATTTTCAGAAATTAACATAAAATCCAGAA  
850 860 870 880 890 900  
GCAAGATTCCGTAAGCTGAGAACTCTGGACAGTTGATCAGCTTTACCTATGGTGCTTTGC  
910 920 930 940 950 960  
CTTTAACTAGAGTGTGTGATGGTAGATTATTTTCAGATATGTATGTAAAACGTTCCTGA  
970 980 990 1000 1010 1020  
ACAATAAGATGTATGAACGGAGCAGAAATAAATACTTTTTCTAATTAATACCTTTAAAAA  
1030  
AAAAAAAAAA

SEQ ID NO:20 (BMS155 protein)

10 20 30 40 50 60  
METLYRVPFLVLECPNLKLKKPPKLLHMPSAMTVYALVVVSYFLITGGIIYDVIVEPPSVG  
70 80 90 100 110 120  
SMTDEHGHQRPVAFLAYRVNGQYIMEGLASSFLFTMGGLGFIIIDRSNAPNIPKLNRFLL  
130 140  
LFIGFVCVLLSFFMARVFMRMKLPGYLMG

## SEQ ID NO:21 (BMS208 cDNA)

10	20	30	40	50	60
GTTGATTGGGTCTAGACCAAAGAACTTTGAGGAACTTGCCCAGAGCCCTGCATGCATCAG					
70	80	90	100	110	120
ACCTACAGCAGACATTGCAGGCCTGAAGAAAGGTGGTCACAAGAGGGGTGGAACATTCCT					
130	140	150	160	170	180
GCAAATGGTTTCAATATATGCAGATGTCTCGATATAGGAATGAAATTACGTCTTTGGAAC					
190	200	210	220	230	240
AACTTAAATAAGTCAAATATACTTGGAGCTTTAAAAATTAAAAGGAGAGAGATTGAGCA					
250	260	270	280	290	300
CCTTTTCTGCTGCCATGACAACCATGCAAGGAATGGAACAGGCCATGCCAGGGGCTGGCC					
310	320	330	340	350	360
CTGGTGTGCCCCAGCTGGGAAACATGGCTGTCATACATTACATCTGTGGAAAGGATTGC					
370	380	390	400	410	420
AAGAGAAGTTCTTGAAGGGAGAACCCAAAGTCCTTGGGGTTGTGCAGATTCTGACTGCCC					
430	440	450	460	470	480
TGATGAGCCTTAGCATGGGAATAACAATGATGTGTATGGCATCTAATACTTATGGAAGTA					
490	500	510	520	530	540
ACCCTATTTCCGTGTATATCGGGTACACAATTTGGGGGTCAGTAATGTTTATTATTTCAG					
550	560	570	580	590	600
GATCCTTGTCAATTGCAGCAGGAATTAGAACTACAAAAGGCCTGGGTCTGGATGGCATGG					
610	620	630	640	650	660
TGCTCCTCTTAAGTGTGCTGGAATTCTGCATTGCTGTGTCCCTCTCTGCCTTTGGATGTA					
670	680	690	700	710	720
AAGTGCTCTGTTGTACCCCTGGTGGGGTTGTGTTAATTCTGCCATCACATTCTCACATGG					

730 740 750 760 770 780  
CAGAAACAGCATCTCCACACCACTTAATGAGGTTTQAGGCCACCAAAGATCAACAGAC

790 800 810 820 830 840  
AAATGCTCCAGAAATCTATGCTGACTGTGACACAAGAGCCTCACATGAGAAATTACCAGT

850 860 870 880 890 900  
ATCCAACCTTCGATACTGATAGACTTGTGATATTATTATTATATGTAATCCAATTATGAA

910 920 930 940 950 960  
CTGTGTGTGTATAGAGAGATAATAAATTCAAAATTATGTTCTCATTTTTTCCCTGGAAC

970 980 990 1000 1010 1020  
TCAATAACTCATTTCCTGGCTCTTTATCGAGAGTACTAGAAGTTAAATTAATAAATAAT

1030 1040 1050 1060 1070 1080  
GCATTTAATGAGGCAACAGCACTTGAAAGTTTTTCATTCATCATAAGAACTTTATATAAA

1090 1100 1110 1120 1130 1140  
GGCATTACATTGGCAAATAAGGTTTGGAAGCAGAAGAGCAAAAAAAGATATTGTTAAAA

1150 1160 1170 1180 1190 1200  
TGAGGCCTCCATGCAAAACACATACTTCCCTCCCATTTATTTAACTTTTTTTTTCTCCT

1210 1220 1230 1240 1250 1260  
ACCTATGGGGACCAAAGTGCTTTTTCCTTCAGGAAGTGGAGATGCATGGCCATCTCCCCC

1270 1280 1290 1300 1310 1320  
TCCCTTTTTCCTTCTCCTGCTTTTCTTCCCATAGAAAGTACCTTGAAGTAGCACAGTC

1330 1340 1350 1360 1370 1380  
CGTCCTTGATGTGCACGAGCTATCATTGAGTAAAAGTATACATGGAGTAAAAATCATA

1390 1400 1410 1420 1430 1440  
TTAAGCATCAGATTCAACTTATATTTTCTATTTTCATCTTCTTCCCTTCCCTTCTCCCACC

1450 1460 1470 1480 1490 1500  
TTCTACTGGGCATAATTATATCTTAATCATATATGGAAATGTGCAACATATGGTATTTGT

1510 1520 1530 1540 1550 1560  
TAAATACGTTTGTGTTTTATTGCAGAGCAAAAATAAATCAAATTAGAAGCAAAAAAAAAAA  
AAA

## SEQ ID NO:22 (BMS208 protein)

10 20 30 40 50 60  
MTTMOGMEQAMPGAGPGVPOLGNMAVIHSHLWKGLOEKFLKGEPKVLGVVOILTALMSLS  
70 80 90 100 110 120  
MGITMMCMASNTYGSNPISVYIGYTIWGSVMFIISGSLIAAGIRTTKGLGLDGMVLLLS  
130 140 150 160  
VLEFCIAVSLSAFGCKVLCCTPGGVVLILPSHSHMAETASPTPLNEV

## SEQ ID NO:23 (BMS235 cDNA)

10 20 30 40 50 60  
CCGGCGGGACGGAGGGCCCGGCAGGAAGATGGGCTCCCGTGGACAGGGACTCTTGCTGGC  
70 80 90 100 110 120  
GTACTGCCTGCTCCTTGCCCTTTGCCTCTGGCCTGGTCCTGAGTCGTGTGCCCCATGTCCA  
130 140 150 160 170 180  
GGGGGAACAGCAGGAGTGGGAGGGGACTGAGGAGCTGCCGTGCGCTCCGGACCATGCCGA  
190 200 210 220 230 240  
GAGGGCTGAAGAACAACATGAAAAATACAGGCCCACTCAGGACCAGGGGCTCCCTGCTTC  
250 260 270 280 290 300  
CCGGTGCTTGCGCTGCTGTGACCCCGGTACCTCCATGTACCCGGCGACCGCCGTGCCCCA  
310 320 330 340 350 360  
GATCAACATCACTATCTTGAAAGGGGAGAAGGGTGACCGGGAGATCGAGGCCTCCAAGG  
370 380 390 400 410 420  
GAAATATGGCAAAACAGGCTCAGCAGGGGCCAGGGGCCACACTGGACCCAAAGGGCAGAA

430 440 450 460 470 480  
GGGCTCCATGGGGGCCCTGGGGAGCGGTGCAAGAGCCACTACGCCGCCCTTTTCGGTGGG  
490 500 510 520 530 540  
CCGGAAGAAGCCCATGCACAGCAACCACTACTACCAGACGGTGATCTTCGACACGGAGTT  
550 560 570 580 590 600  
CGTGAACCTCTACGACCACTTCAACATGTTACCGGCAAGTTCTACTGCTACGTGCCCGG  
610 620 630 640 650 660  
CCTCTACTTCTTCAGCCTCAACGTGCACACCTGGAACCAGAAGGAGACCTACCTGCACAT  
670 680 690 700 710 720  
CATGAAGAACGAGGAGGAGGTGGTGATCTTGTTCGCGCAGGTGGGCGACCGCAGCATCAT  
730 740 750 760 770 780  
GCAAAGCCAGAGCCTGATGCTGGAGCTGCGAGAGCAGGACCAGGTGTGGGTACGCCTCTA  
790 800 810 820 830 840  
CAAGGGCGAACGTGAGAACGCCATCTTCAGCGAGGAGCTGGACACCTACATCACCTTCAG  
850 860 870 880 890 900  
TGGCTACCTGGTCAAGCACGCCACCGAGCCCTAGCTGGCCGGCCACCTCCTTTCTCTCG  
910 920 930 940 950 960  
CCACCTTCCACCCCTGCGCTGTGCTGACCCACCGCCTCTTCCCCGATCCCTGGACTCCG  
970 980 990 1000 1010 1020  
ACTCCCTGGCTTTGGCATTCACTGAGACGCCCTGCACACACAGAAAGCCAAAGCGATCGG  
1030 1040 1050 1060 1070 1080  
TGCTCCAGATCCCGCAGCCTCTGGAGAGAGCTGACGGCAGATGAAATCACCAGGGCGGG  
1090 1100 1110 1120 1130 1140  
GCACCCGCGAGAACCCTCTGGGACCTTCCGCGGCCCTCTCTGCACACATCCTCAAGTGAC  
1150 1160 1170 1180 1190 1200  
CCCGCACGGCGAGACGCGGTGGCGGCAGGGCGTCCCAGGGTGCGGCACCGCGGCTCCAG  
1210 1220 1230 1240 1250 1260

TCCTTGGAATAATTAGGCAAATTCTAAAGGTCTCAAAGGAGCAAAGTAAACCGTGGAG  
1270 1280 1290 1300 1310 1320  
GACAAAGAAAAGGGTTGTTATTTTTGTCTTTCCAGCCAGCCTGCTGGCTCCCAAGAGAGA  
1330 1340 1350 1360 1370 1380  
GGCCTTTTCAGTTGAGACTCTGCTTAAGAGAAGATCCAAAGTTAAAGCTCTGGGGTCAGG  
1390 1400 1410 1420 1430 1440  
GGAGGGGCGGGGGCAGGAACTACCTCTGGCTTAATTCTTTTAAGCCACGTAGGAACCTT  
1450 1460 1470 1480 1490 1500  
TCTTGAGGGATAGGTGGACCCTGACATCCCTGTGGCCTTGCCCAAGGGCTCTGCTGGTCT  
1510 1520 1530 1540 1550 1560  
TTCTGAGTCACAGCTGCGAGGTGATGGGGGCTGGGGCCCCAGGCGTCAGCCTCCCAGAGG  
1570 1580 1590 1600 1610 1620  
GACAGCTGAGCCCCCTGCCTTGGCTCCAGGTTGGTAGAAGCAGCCGAAGGGCTCCTGACA  
1630 1640 1650 1660 1670 1680  
GTGGCCAGGGACCCCTGGGTCCCCCAGGCCTGCAGATGTTTCTATGAGGGGCAGAGCTCC  
1690 1700 1710 1720 1730 1740  
TGGTACATCCATGTGTGGCTCTGCTCCACCCCTGTGCCACCCCAGAGCCCTGGGGGGTGG  
1750 1760 1770 1780 1790 1800  
TCTCCATGCCTGCCACCCTGGCATCGGCTTTCTGTGCCGCCTCCACACAAATCAGCCCC  
1810 1820 1830 1840 1850 1860  
AGAAGGCCCCGGGGCCTTGGCTTCTGTTTTTTATAAAACACCTCAAGCAGCACTGCAGTC  
1870 1880 1890 1900 1910 1920  
TCCCATCTCCTCGTGGGCTAAGCATCACCGCTTCCACGTGTGTTGTGTTGGTTGGCAGCA  
1930 1940 1950 1960 1970 1980  
AGGCTGATCCAGACCCCTTCTGCCCCCACTGCCCTCATCCAGGCCTCTGACCAGTAGCCT  
1990 2000 2010 2020 2030 2040  
GAGAGGGGCTTTTTCTAGGCTTCAGAGCAGGGGAGAGCTGGAAGGGGCTAGAAAGCTCCC



2050      2060      2070      2080      2090      2100  
 GCTTGTCTGTTTCTCAGGCTCCTGTGAGCCTCAGTCCTGAGACCAGAGTCAAGAGGAAGT  
 2110      2120      2130      2140      2150      2160  
 ACACGTCCCAATCACCCGTGTCAGGATTCACTCTCAGGAGCTGGGTGGCAGGAGAGGCAA  
 2170      2180      2190      2200      2210      2220  
 TAGCCCCTGTGGCAATTGCAGGACCAGCTGGAGCAGGGTTGCGGTGTCTCCACGGTGCTC  
 2230      2240      2250      2260      2270      2280  
 TCGCCCTGCCCATGGCCACCCCAGACTCTGATCTCCAGGAACCCCATAGCCCCTCTCCAC  
 2290      2300      2310      2320      2330      2340  
 CTCACCCCATGTTGATGCCCAGGGTCACTCTTGCTACCCGCTGGGCCCCCAAACCCCGC  
 2350      2360      2370      2380      2390      2400  
 TGCCTCTCTTCTTCCCCCATCCCCACCTGGTTTTGACTAATCCTGCTTCCCTCTCTG  
 2410      2420      2430      2440      2450      2460  
 GGCCTGGCTGCCGGGATCTGGGGTCCCTAAGTCCCTCTCTTTAAGAACTTCTGCGGGTC  
 2470      2480      2490      2500      2510      2520  
 AGACTCTGAAGCCGAGTTGCTGTGGGCGTGCCCGGAAGCAGAGCGCCACACTCGCTGCTT  
 2530      2540      2550      2560      2570      2580  
 AAGCTCCCCCAGCTCTTTCCAGAAAACATTAAACTCAGAATTGTGTTTTCAAAAAAAAAA  
 2590  
 AAAAAAAAAA

SEQ ID NO:24 (BMS235 protein)

10      20      30      40      50      60  
 MGSRGOGLLLAYCLLLAFASGLVLSRVPHVQGEQQEWEGTEELPSPPDHAERAEQHEKY  
 70      80      90      100      110      120  
 RPSQDQGLPASRCLRCCDPGTSHPATAVPQINITILKEKGDRGDRGLQKRYGKTGSAG  
 130      140      150      160      170      180  
 ARGHTGPKGQKGSMSGAPGERCKSHYAAPSVGRKKPMHSNHYYQTVIFDTEFVNLVDHFNH

190 200 210 220 230 240  
FTGKFYCYVPGLYFFSLNVHTWNQKETYLHIMKNEEEVVILFAQVGDRSIMQSLSMLEL

250 260 270 280  
REQDQVWVRLYKGERENAI FSEELDTYITFSGYLVKHATEP

SEQ ID NO:25 (BMS240 cDNA)

10 20 30 40 50 60  
GGACTTGAGCGAGCCAGTTGCCGATTATTCTATTTCCCTCCCTCTCTCCCGCCCCGTA

70 80 90 100 110 120  
TCTCTTTTCACCTTCTCCACCCCTCGCTCGCGTAGCCATGGCGGAGCCGTCGGCGGCCA

130 140 150 160 170 180  
CTCAGTCCCATTCCATCTCCTCGTCGTCCTTCGGAGCCGAGCCGTCGCGCCCGGCGGCG

190 200 210 220 230 240  
GCGGGAGCCCAGGAGCCTGCCCCGCCCTGGGGACGAAGAGCTGCAGCTCCTCCTGTGCGG

250 260 270 280 290 300  
TGCACGATCTGATTTTCTGGAGAGATGTGAAGAAGACTGGGTTTGTCTTTGGCACCACGC

310 320 330 340 350 360  
TGATCATGCTGCTTTCCTGGCAGCTTTCAGTGTTCATCAGTGTGGTTTCTTACCTCATCC

370 380 390 400 410 420  
TGGCTCTTCTCTCTGTCAACATCAGCTTCAGGATCTACAAGTCCGTCATCCAAGCTGTAC

430 440 450 460 470 480  
AGAAGTCAGAAGAAGGCCATCCATTCAAAGCCTACCTGGACGTAGACATTACTCTGTCTCT

490 500 510 520 530 540  
CAGAAGCTTCCATAATTACATGAATGCTGCCATGGTGACATCAACAGGGCCCTGAAAC

550 560 570 580 590 600  
TCATTATTCTCTCTTTCTGGTAGAAGATCTGGTTGACTCCTTGAAGCTGGCTGTCTTCA

610 620 630 640 650 660  
TGTGGCTGATGACCTATGTTGGTGCTGTTTTTAACGGAATCACCCCTTCTAATTCTTGCTG

670 680 690 700 710 720  
AACTGCTCATTTCAGTGTCCCGATTGTCTATGAGAAGTACAAGACCCAGATTGATCACT

730 740 750 760 770 780  
ATGTTGGCATCGCCCGAGATCAGACCAAGTCAATTGTTGAAAAGATCCAAGCAAACTCC

790 800 810 820 830 840  
CTGGAATCGCCAAAAAAGGCAGAAATAAGTACATGGAAACCAGAAATGCAACAGTTACT

850 860 870 880 890 900  
AAAACACCATTTAATAGTTATAACGTCGTTACTTGTACTATGAAGGAAAATACTCAGTGT

910 920 930 940 950 960  
CAGCTTGAGCCTGCATTCCAAGCTTTTTTTTTTAATTGTTGTTTCTCCCATCCTTTCCC

970 980 990 1000 1010 1020  
TTTAACCCCTCAGTATCAAGCACAAAAATTGATGGACTGATAAAAGAACTATCTTAGAACT

1030 1040 1050 1060 1070 1080  
CAGAAGAAGAAAGAAATCAAATTCATAGGATAAGTCAATACCTTAATGGTGGTAGAGCCTT

1090 1100 1110 1120 1130 1140  
TACCTGTAGCTTGAAAGGGGAAAGATTGGAGGTAAGAGAGAAAATGAAAGAACACCTCTG

1150 1160 1170 1180 1190 1200  
GGTCCTTCTGTCCAGTTTTTCAGCACTAGTCTTACTCAGCTATCCATTATAGTTTTGCCCT

1210 1220 1230 1240 1250 1260  
TAAGAAGTCATGATTAACCTTATGAAAAAATTATTTGGGGACAGGAGTGTGATACCTTCCT

1270 1280 1290 1300 1310 1320  
TGGTTTTTTTTTGCAGCCCTCAAATCCTATCTTCCTGCCCCACAATGTGAGCAGCTACCC

1330 1340 1350 1360 1370 1380  
CTGATACTCCTTTTCTTTAATGATTTAACTATCAACTTGATAAATAACTTATAGGTGATA

1390 1400 1410 1420 1430 1440  
GTGATAATTCCTGATTCCAAGAATGCCATCTGATAAAAAAGAATAGAAATGGAAAGTGGG  
1450 1460 1470 1480 1490 1500  
ACTGAGAGGGAGTCAGCAGGCATGCTGCGGTGGCGGTCACTCCCTCTGCCACTATCCCCA  
1510 1520 1530 1540 1550 1560  
GGGAAGGAAAGGCTCCGCCATTGGGAAAGTGGTTTCTACGTCACTGGACACCGGTTCTG  
1570 1580 1590 1600 1610 1620  
AGCATTAGTTTGAGAACTCGTTCCCGAATGTGCTTTCCTCCCTCTCCCCTGCCACCTCA  
1630 1640 1650 1660  
AGTTTAAATAAATAAGGTTGTACTTTTCTACTATAAAAAAAAAAAAAA

SEQ ID NO:26 (BMS240 protein)

10 20 30 40 50 60  
MAEPSAATQSHSISSSSFGAEPSPAGGGSPGACPALGTKSCSSSCAVHDLIFWRDVKKT  
70 80 90 100 110 120  
GFVFGTTLIMLLSLAAFVISVVSYLILALLSVTISFRIYKSVIQAVQKSEEGHPFKAYL  
130 140 150 160 170 180  
DVDITLSSEAFHNYMNAAMVHINRAKLIIRLFLVEDLVDSLKLAVFMWLMITYVGAVFNG  
190 200 210 220 230  
ITLLILAELLIFSVPIVYEKYKTQIDHYVGIARDQTKSIVEKIQAKLPGIAKKAE

## Sequence of BMS53 cDNA (Range: 1 to 1697)

Seq ID NO:27

```

      10      20      30      40      50      60
70
CTTCATCCTGCCCGCGTCACTGAGAGGATGTTCAACCAGAATGTGGTGGCCCAGCTCTGGTACTTCGTG

      80      90     100     110     120     130
140
AAGTGCATCTACTTCGCCCTGTCCGCCTACCAGATCCGCTGCGGCTACCCACCCGCATCCTCGGCAACT

      150     160     170     180     190     200
210
TCCTCACCAAGAAGTACAATCATCTCAACCTCTTCTCTTCCAGGGGTTCGGGCTGGTGCCGTTCTTGGT

      220     230     240     250     260     270
280
GGAGCTGCGGCAGTGATGGACTGGGTGTGGACGGACACCACGCTGTCCCTGTCCAGCTGGATGTGTGTG

      290     300     310     320     330     340
350
GAGGACATCTATGCCAACATCTTCATCATCAAATGCAGCCGAGAGACAGAGAAGAAATACCCGAGCCCA

      360     370     380     390     400     410
420
AAGGGCAGAAGAAGAAGATCGTCAAGTACGGCATGGGTGGCCTCATCATCCTCTTCTCATCGCCAT

      430     440     450     460     470     480
490
CATCTGGTTCCCGCTGCTCTTCATGTCGCTGGTGCCTCCGTGGTTGGGGTTGTCAACCAGCCCATCGAT

      500     510     520     530     540     550
560
GTCACCGTCACCCTGAAGCTGGGCGGCTATGAGCCGCTGTTACCATGAGCGCCAGCAGCCGTCCATCA

      570     580     590     600     610     620
630
TCCCCTTCACGGECCAGGCCTATGAGGAGCTGTCCCGGCAGTTTGACCCCCAGCCGCTGGCCATGCAGTT

      640     650     660     670     680     690
700
CATCAGCCAGTACAGCCCTGAGGACGTCGTACGGCGCAGATTGAGGGCAGCTCCGGGGCGCTGTGGCGC

      710     720     730     740     750     760

```

770

ATCAGTCCCCCAGCCGTGCCCAGATGAAGCGGGAGCTCTACAACGGCACGGCCGACATCACCTGCGCT

780 790 800 810 820 830

840

TCACCTGGAACCTCCAGAGGGACCTGGCGAAGGGAGGCACTGTGGAGTATGCCAACGAGAAGCACATGCT

850 860 870 880 890 900

910

GGCCCTGGCCCCAACAGCACTGCACGGCGGCAGCTGGCCAGCCTGCTCGAGGGCACCTCGGACCAGTCT

920 930 940 950 960 970

980

GTGGTCATCCCAATCTCTTCCCAAGTACATCCGTGCCCCAACGGGCCGAAGCCAACCTGTGAAGC

990 1000 1010 1020 1030 1040

1050

AGCTGCAGCCCAATGAGGAGGCCGACTACCTCGGCGTGCGTATCCAGCTGCGGAGGGAGCAGGGTGCGGG

1060 1070 1080 1090 1100 1110

1120

GGCCACCGGCTTCCTCGAATGGTGGGTCACTCGAGCTGCAGGAGTGCCGACCGACTGCAACCTGTGCCC

1130 1140 1150 1160 1170 1180

1190

ATGGTCATTTTCAGTGACAAGGTCAGCCACCGAGCCTCGGCTTCCTGGCTGGCTACGGCATCATGGGGC

1200 1210 1220 1230 1240 1250

1260

TGTAAGTGTCCATCGTGCTGGTCACTCGGCAAGTTGCTGCGCGGATTCTTCAGCGAGATCTCGCACTCCAT

1270 1280 1290 1300 1310 1320

1330

TATGTTGAGGAGCTGCCGTGCGTGGACCGCATCCTCAAGCTCTGCCAGGACATCTTCCTGGTGCAGGAG

1340 1350 1360 1370 1380 1390

1400

ACTCGGAGCTGGAGCTGGAGGAGGAGTTGTACGCCAAGCTCATCTTCCTCTACCGCTCACCGGAGACCA

1410 1420 1430 1440 1450 1460

1470

TGATCAAGTGGACTCGTGAGAAGGAGTAGGAGCTGCTGCTGGCGCCGAGAGGGAAGGAGCCGGCCTGCT

1480 1490 1500 1510 1520 1530

1540

GGGCAGCGTGCCACAAGGGGCGGCACTCCTCAGGCCGGGGAGCCACTGCCCCGTCCAAGGCCGCCAGC

1550 1560 1570 1580 1590 1600  
1610

TGTGATGCATCCTCCCGGCCTGCCTGAGCCCTGATGCTGCTGTCAGAGAAGGACACTGCGTCCCCACGGC

1620 1630 1640 1650 1660 1670  
1680

CTGCGTGGCGCTGCCGTCCCCACGTGTACTGTAGAGTTTTTTTTTAATTAAAAATGTTTTATTATA

1690  
CAAAAAAAAAAAAAAA

Seq ID NO: 28

### Sequence of the predicted BMS53 polypeptide (Range: 1 to 466)

10 20 30 40 50 60  
70  
MFNQNVVAQLWYFVKCIYFALSAYQIRCGYPTRILGNFLTKKYNHLNLFQGFRLVPPFLVELRAVMDWV

80 90 100 110 120 130  
140  
WTDTTLSLSSWMCVEDIYANIFIICKSRETEKKYPQPKGQKKKIVKYGMGGLIILFLIAIIWFPLLFMS

150 160 170 180 190 200  
210  
LVRSVVGVVNQPIDVTTLKLGGEPLFTMSAQQPSIIPFTAQAYEELSROQFDPQPLAMQFISQYSPEDV

220 230 240 250 260 270  
280  
VTAQIEGSSGALWRISPPSRAQMKRELYNGTADITLRFWNFQDLAKGGTVEYANEKHMALAPNSTAR

290 300 310 320 330 340  
350  
RQLASLLEGTSDQSVVIPNLFPKYIRAPNGPEANPVKQLQPNEEADYLGVRIQLRREQGAGATGFLEWWV

360 370 380 390 400 410  
420  
IELQECRTDCNLLPMVIFSDKVSPPSLGFLAGYGIMGLYVSIVLVIGKFVRGFFSEISHSIMFEELPCVD

430 440 450 460  
RILKLCQDIFLVRETRERELEEELYAKLIFLYRSPETMIKW TREKE

Sequence of BMS100 cDNA (Range: 1 to 1333) *Seq ID NO: 29*

```

      10      20      30      40      50      60
GGTGGGTGCATCCTGCGCTGCGGCGGGCGCGCTACCCAGACGCTGGTGTGCAGAGCCACA

      70      80      90     100     110     120
TGAAGCCTGCTGGGGACTGGGGGCCAGGGAGCAGCAAGCCAGCTGGGACTGAGGCGGACG

     130     140     150     160     170     180
CTGTCTCAGGGAGACGCTGACTCGCAAAGACACTCCCTTCCTTGTGCCTGGGTAAAAAGT

     190     200     210     220     230     240
CTCCTCCTGGGGTCCCTGGCCATCCTGAATATCCAGAATGGTGTTCCTGAAGTTCCTCTG

     250     260     270     280     290     300
CATGAGTTTCTTCTGCCACCTGTGTCAAGGCTACTTCGATGGCCCCCTCTACCCAGAGAT

     310     320     330     340     350     360
GTCCAATGGGACTCTGCACCACTACTTCGTGCCCGATGGGGACTATGAGGAGAACGATGA

     370     380     390     400     410     420
CCCCGAGAAGTGCCAGCTGCTCTTCAGGGTGAGTGACCACAGGCGCTGCTCCCAGGGGGA

     430     440     450     460     470     480
GGGGAGCCAGGTTGGCAGCCTGCTGAGCCTCACCTGCGGGAGGAGTTCACCGTGCTGGG

     490     500     510     520     530     540
CCGCCAGGTGGAGGATGCTGGGCGCGTGCTGGAGGGCATCAGCAAAAGCATCTCCTACGA

     550     560     570     580     590     600
CCTAGACGGGGAAGAGAGCTATGGCAAGTACCTGCGGCGGGAGTCCCACCAGATCGGGGA

     610     620     630     640     650     660
TGCCTACTCCAACCTCGGACAAATCCCTCACTGAGCTGGAGAGCAAGTTCAAGCAGGGCCA

     670     680     690     700     710     720
GGAACAGGACAGCCGGCAGGAGAGCAGGCTCAACGAGGACTTCTGGGAATGCTGGTCCA

     730     740     750     760     770     780
CACCAGGTCCCTGCTGAAGGAGACACTGGACATCTCTGTGGGGCTCAGGGACAAATACGA

     790     800     810     820     830     840
GCTGCTGGCCCTCACCATTAGGAGCCATGGGACCCGACTAGGTCCGCTGAAAAATGATTA

     850     860     870     880     890     900
TCTTAAAGTATAGGTGGAAGGATACAAATGCTAGAAAGAGGGAATCAAATCAGCCCCGTT

     910     920     930     940     950     960
TTGGAGGGTGGGGGACAGAAGATGGGGCTACATTTCCCCCATACCTACTATTTTTTTATA

     970     980     990    1000    1010    1020
TCCCGATTTGCACTTTGAGAATACATCTAAGGTCATCTTTCAAAGAGAAAAATGGACA

```



1030 1040 1050 1060 1070 1080  
 CTTGAGTGACTTTGTTTTAGTTTTGTTTTGTACATTATTTATGTGATTGTTATGGAAT  
 1090 1100 1110 1120 1130 1140  
 TGTACACCTGGAAAGAACAATTTTAAGCAATGTCATTTCTAGATGGGTTTCTAATTCTGCA  
 1150 1160 1170 1180 1190 1200  
 GAGACACCCGTTTCAGCCACATCTAAAAGAGCACAGTTTATGTGGTGCGGAATTAAACTT  
 1210 1220 1230 1240 1250 1260  
 CCCCATCCTGCAGATTATGTGGAAATACCCAAAGATAATAGTGCATAGCTCCTTTTCAGCC  
 1270 1280 1290 1300 1310 1320  
 TCTAGCCTTCACTCCTGGGCTCCAAAAGCTATCCAGTTGCCTGTTTTTCAAATGAGGTT  
 1330  
 CAAGGTGCTGCTT

211) Sequence of the predicted BMS100 polypeptide (Range: 1 to 520) ID NO: 3

10 20 30 40 50 60  
MVELKEFCMSFECHLCQGYFDGPLYPEMSNGTLHHYFVPDGDYEENDDPEKCQLLFRVSD  
 70 80 90 100 110 120  
 HRRCSQGEQSQVGSLLSLTLREEFTVLGRQVEDAGRVLEGISKISISYDLGDEESYGKYL  
 130 140 150 160 170 180  
 RESHQIGDAYSNSDKSLTELESKFQEQEQDSRQESRLNEDFLGMLVHTRSLKETLDIS  
 190 200 210  
 VGLRDKYELLALTIRSHGTRLGRLKNDYLKV

## Sequence of BMS199 cDNA (Range: 1 to 1102)

SEQ ID NO: 31

```

      10      20      30      40      50      60
GTCTTGGGGTCCCTGGCTGGGTGGCCAGACCCCGAAGCCAGCGCTGGGAAGGGCTGCGGA

      70      80      90     100     110     120
TGCCCGGGTCAGAGGAAGGGCAGGTCCAAGGACACGCGGGTCTGGTCCTGGGCAAGAAC

     130     140     150     160     170     180
CGCCCCCTCTCCGGGCCTGCTTCAGTCTTCCTTTGCAGAACACGGGCCAGGCCCTTCC

     190     200     210     220     230     240
CTCTGCCCCCGGGTGCTTGAAGTCTAGCCCCATCCTGGTCCAATGCGCTCTTGGTAGCCT

     250     260     270     280     290     300
CCTTTCCAGCTGCCCCGCCCGCGCCATGCGCCCTTACTGCCCCCTGCGCCTGTGCCGGC

     310     320     330     340     350     360
TGTGGCCCCCGCAACCTCCCTCCCGGCTCCTCGGAGCGGCCCGCGGCAGCGGTCCAGAC

     370     380     390     400     410     420
CCAGTACTTATTATGAAGTGTGGGGGTGCATCCTGGTGCCAGCACTGAGGAAGTTAAAC

     430     440     450     460     470     480
GAGCTTTCCTCTCCAAGTCCAAAGAGCTGCACCCAGACCGGGACCCTGGGAACCCAAGCC

     490     500     510     520     530     540
TGACACAGCCGCTTTGTGGAGCTGAGCGAGGCATACCGTGTGCTCAGCCGTGAGCAGAGCC

     550     560     570     580     590     600
GCCGCAGCTATGATGACCAGCTCCGCTCAGGTAGTCCCCCAAAGTCTCCACGAACCACAG

     610     620     630     640     650     660
TCCATGACAAGTCTGCCCACCAAACACACAGCTCCTGGACACCCCCCAACGCACAGTACT

     670     680     690     700     710     720
GGTCCCAGTTTCACAGCGTGAGGCCACAGGGGCCCCAGTTGAGGCAGCAGCAACACAAAC

     730     740     750     760     770     780
AAAACAAACAAGTGCTGGGGTACTGCCTCCTCCTCATGCTGGCGGGCATGGGCCTGCACT

     790     800     810     820     830     840
ACATTGCCTTCAGGAAGGTGAAGCAGATGCACCTTAACCTTCATGGATGAAAAGGATCGGA

     850     860     870     880     890     900
TCATCACAGCCTTCTACAACGAAGCCCGGGCACGGGCCAGGGCCAACAGAGGCATCCTTC

     910     920     930     940     950     960
AGCAGGAGCGACAACGGCTAGGGCAGCGGCAGCCGCCACCATCCGAGCCAACCCAAGGCC

     970     980     990    1000    1010    1020
CCGAGATCGTGCCCCGGGGCGCCGGCCCCTGAGGGGCTCACCTGGATGGGGCCTGCAGTG

```

1030 1040 1050 1060 1070 1080  
CGTTCCCGCTTTGCTTCCTTCCTGGACGGCCCGCTCCCCGAAACGCGCGCAATAAAGTG  
1090 1100  
ATTTCGAGAAAAAAAAAAAAA

241) Sequence of the predicted BMS199 polypeptide (Range: 1 to SEQ ID NO:32

10 20 30 40 50 60  
MPPLLPLRLCRLWPRNPPSRLLGAAAGORSRPSTYYELLGVHPGASTE EVKRAFFSKSKE  
70 80 90 100 110 120  
LHPDRDPGNPSLHSRFVELSEAYRVLSREQSRRSYDDQLRSGSPPKSPRTTVHDKSAHOT  
130 140 150 160 170 180  
HSSWTPPNAQYWSQFHSVRPQGPQLRQQQHKQNKQVLGYCLLLMLAGMGLHYIAFRKVKQ  
190 200 210 220 230 240  
MHLNFMDEKDRIITAFYNEARARARANRGILQQERQRLGQRQPPPPSEPTQGPEIVPRGAG

P

Seq ID NO:33

## Sequence of BMS206 cDNA (Range: 1 t 966)

```

      10      20      30      40      50      60
GAGAAGCATCGAGGCTATAGGACGCAGCTGTTGCCATGACGGCCCAGGGGGCCTGGTGG

      70      80      90     100     110     120
CTAACCGAGGCCGCGCTTCAAGTGGGCCATTGAGCTAAGCGGGCCTGGAGGAGGCAGCA

      130     140     150     160     170     180
GGGGTCGAAGTGACCGGGGCAGTGGCCAGGGAGACTCGCTCTACCCAGTGGTTACTTGG

      190     200     210     220     230     240
ACAAGCAAGTGCCGTGATACCAGCGTGCAAGAGACAGACCGGATCCTGGTGGAGAAGCGCT

      250     260     270     280     290     300
GCTGGGACATCGCCTTGGGTCCCCTCAAACAGATTCCCATGAATCTCTTCATCATGTACA

      310     320     330     340     350     360
TGGCAGGCAATACTATCTCCATCTTCCCTACTATGATGGTGTGTATGATGGCCTGGCGAC

      370     380     390     400     410     420
CCATTTCAGGCACTTATGGCCATTTTCAGCCACTTTCAAGATGTTAGAAAGTTCAAGCCAGA

      430     440     450     460     470     480
AGTTTCTTCAGGGTTTGGTCTATCTCATTTGGGAACCTGATGGGTTTGGCATTTGGCTGTTT

      490     500     510     520     530     540
ACAAGTGCCAGTCCATGGGACTGTTACCTACACATGCATCGGATTGGTTAGCCTTCATTG

      550     560     570     580     590     600
AGCCCCCTGAGAGAATGGAGTTTCAGTGGTGGAGGACTGCTTTTGTGAACATGAGAAAGCA

      610     620     630     640     650     660
GCGCCTGGTCCCCTATGTATTTGGGTCTTATTTACATCCTTCTTTAAGCCCAGTGGCTCCT

      670     680     690     700     710     720
CAGCATACTCTTAAACTAATCACTTATGTTAAAAAGAACCAAAGACTCTTTTCTCCATG

      730     740     750     760     770     780
GTGGGGTGACAGGTCCCTAGAAGGACAAATGTGCATATTACGACAAACACAAAGAACTATA

      790     800     810     820     830     840
CCATAACCCAAGGCTGAAAATAATGTAGAAAACCTTATTTTGTTCAGTACAGAGCAA

      850     860     870     880     890     900
AACAACAACAAAAAACATAACTATGTAAACAAGAGAATAACTGCTGCTAAATCAAGAAC

      910     920     930     940     950     960
TGTTGCAGCATCTCCTTTCAATAAATTAAATGGTTGAGAACAATGCATAAAAAAAAAA

AAAAAA
```

~~SEQ~~ ID NO: 34

Sequence of the predicted BMS206 polypeptide (Range: 1 to 183)

10 20 30 40 50 60  
MTAQGGLVANRGRFRKWAIELSGPGGSRGRSDRGSGQGDSLYPVGYLDKQVPDTSVQET  
70 80 90 100 110 120  
DRILVEKRCWDIALGPLKQIPMNLFIYMAGNTISIFPTMMVCMMAWRPIQALMAISATF  
130 140 150 160 170 180  
KMLESSSQKFLQGLVYLIGNLMGLALAVYKCQSMGLLPTHASDWLAFIEPPERMEFSGGG  
LLL

Seq ID NO: 35

## Sequence of BMS242 cDNA (Range: 1 t 1570)

```

      10      20      30      40      50      60
GGGCCGGGCGCGGCGCAGAGGCGGGCGCCTACCAGCCGGCAGCTCCGGAGCTGCCCGCGC

      70      80      90     100     110     120
CATGTCCGCGCACAAATCGGGGCACCGAGCTCGACCTTAGCTGGATCTCCAAAATACAAGT

      130     140     150     160     170     180
GAATCACCCGGCAGTTCTGAGGCGTGCAGAACAAATCCAGGCTCGCAGAACCGTGAAAAA

      190     200     210     220     230     240
GGAGTGGCAGGCTGCTTGGCTCCTGAAAGCTGTACCTTTATAGATCTTACTACACTTTC

      250     260     270     280     290     300
AGGTGATGATACATCTTCCAACATTCAAAGGCTCTGTTATAAAGCCAAATACCCAATCCG

      310     320     330     340     350     360
GGAAGATCTCTTAAAAGCTTTAAATATGCATGATAAAGGCATTACTACAGCCGCCGTTTG

      370     380     390     400     410     420
TGTTTATCCCGCCCGGGTGTGTGATGCTGTAAAAGCACTCAAGGCTGCAGGCTGTAATAT

      430     440     450     460     470     480
CCCTGTGGCATCAGTGGCCGCTGGATTTCAGCTGGACAGACTCATTGGAAGACACGATT

      490     500     510     520     530     540
AGAAGAGATCAGATTGGCTGTGGAAGATGGAGCTACAGAAATCGACGTGGTAATTAACAG

      550     560     570     580     590     600
AAGCTTGGTGCTGACAGGCCAGTGGGAAGCCCTGTACGATGAGATTTCGTCACTTCGCAA

      610     620     630     640     650     660
GGCCTGTGGGGAGGCTCATCTTAAAACTATATTAGCGACAGGAGAACTTGGAACTCTTAC

      670     680     690     700     710     720
TAATGTCTATAAAGCCAGTATGATAGCAATGATGGCAGGATCAGATTTTATTAAGACCTC

      730     740     750     760     770     780
TACTGGAAAAGAAACAGTAAATGCCACCTTCCCGGTAGCTATAGTAATGCTGCGGGCCAT

      790     800     810     820     830     840
TAGAGATTTCTTCTGGAAAACCTGGAAACAAGATAGGGTTTAAACCAGCAGGAGGCATCCG

      850     860     870     880     890     900
CAGTGCAAAGGATTCCCTTGCTTGGCTCTCTCTTGTAAAGGAGGAGCTTGGAGATGAGTG

      910     920     930     940     950     960
GCTGAAGCCAGAACTCTTTTGAATAGGTGCCAGTACTCTGCTCTCGACATTGAGAGGCA

      970     980     990    1000    1010    1020
GATTTACCATCATGTGACTGGAAGATATGCAGCTTATCATGATCTTCCAATGTCTTAAAT

```

```

      1030      1040      1050      1060      1070      1080
CAGTCACCAGTTCAGAAAAGTTCTTTACGACAATGTTTAAAAATTATTTTCTACGTAA

      1090      1100      1110      1120      1130      1140
TTGCTAAAATTATTTAATTAAAAAATTGGGCAGTAGGTAAGTGGCATTCTCTCTTTAAA

      1150      1160      1170      1180      1190      1200
ATTTCTACCGAACTTAATGGAATGGAAAAAGCAAATCATCCACATGTGGTACTCATTTTC

      1210      1220      1230      1240      1250      1260
AGGCACATCTGAAATGATCTTAATTACTAGAAGATCTGCACTATTAACCTTTGTGAAGAGT

      1270      1280      1290      1300      1310      1320
TTCTCTAAAAAAGTTTAAGTAAAATGTTAATGGTAGCTTTGATAACATCAAATTTCTAAGG

      1330      1340      1350      1360      1370      1380
GAGAAAAAACAATATTAAACCGCCCAAGCAGTGTGCCCTAGCAGAGGAAAATGCAACAT

      1390      1400      1410      1420      1430      1440
CTCGCAAGCGCTGCTGTAACGACTTCAGGAGTCACTGATTGAGCACTAATTTCTGCTGT

      1450      1460      1470      1480      1490      1500
GAAAACTCATCTTTTCATTTTGGCGTGGATAGGCGCTTTTATTAATGTTGTCTTAATGA

      1510      1520      1530      1540      1550      1560
AATTTCTGACATTGTTCATATACAACGATGAATATCATTAATAATTTTAAAAATAAAAAAA

      1570
AAAAAAAAAA

```

Seq ID  
No: 36

Sequence of the predicted BMS242 polypeptide (Range: 1 to 318)

```

      10      20      30      40      50      60
MSAHNRGTDLDSWISKIQVNHPAVLRRAEQIQARRTVKKEWQAALLKAVTFIDLTTLS

      70      80      90      100     110     120
GDDTSSNIQRLCYKAKYPIREDLLKALNMHDKGITTAAVCVYPARVCDVKAALKAAGCNI

      130     140     150     160     170     180
PVASVAAGFPAGQTHLKTREEIRLAVEDGATEIDVVINRSLVLTGQWEALYDEIRQFRK

      190     200     210     220     230     240
ACGEAHLKTI LATGELGTLTNVYKASMIAMMAGSDFIKTSTGKETVNATFPVAIVMLRAI

      250     260     270     280     290     300
RDFFWKTGNKIGFKPAGGIRSAKDSLAWLSLVKEELGDEWLKPELFRIGASTLLSDIERQ

      310
IYHHVTGRYAAAYHDL PMS

```

SEQ ID NO: 37

Sequence of BMS37 cDNA (Range: 1 to 1542)

```

      10      20      30      40      50      60
CCAACCTCCAACCTCCCTGTCCTGTCTAGGTAACCCCTCCACCCCGCCATTCTCCTATCC

      70      80      90     100     110     120
CGTGTCTGTCCCCATCCCTGTGACCCCTGACCCCTGGCCTTTGCCACTCCCCAGGGACCG

     130     140     150     160     170     180
ATGATGTGGCGACCATCAGTTCTGCTGCTTCTGTTGCTACTGAGGCACGGGGCCCAGGGG

     190     200     210     220     - 230     240
AAGCCATCCCCAGACGCAGGCCCTCATGGCCAGGGGAGGGTGCACCAGGCGGCCCCCTG

     250     260     270     280     290     300
AGCGACGCTCCCCATGATGACGCCCACGGGAACCTCCAGTACGACCATGAGGCTTTCCTG

     310     320     330     340     350     360
GGACGGGAAGTGGCCAAGGAATTCGACCAACTCACCCAGAGGAAAGCCAGGCCCGTCTG

     370     380     390     400     410     420
GGGCGGATCGTGGACCGCATGGACCGCGCGGGGGACGGCGACGGCTGGGTGTGCTGGCC

     430     440     450     460     470     480
GAGCTTCGCGCGTGGATCGCGCACACGCAGCAGCGGCACATACGGGACTCGGTGAGCGCG

     490     500     510     520     530     540
GCCTGGGACACGTACGACACGGACCGCGACGGGCGTGTGGGTGGGAGGAGCTGCGCAAC

     550     560     570     580     590     600
GCCACCTATGGCCACTACGCGCCCCGGTGAAGAATTTTCATGACGTGGAGGATGCAGAGACC

     610     620     630     640     650     660
TACAAAAAGATGCTGGCTCGGGACGAGCGGCGTTTCCGGGTGGCCGACCAGGATGGGGAC

     670     680     690     700     710     720
TCGATGGCCACTCGAGAGGAGCTGACAGCCTTCCTGCACCCCGAGGAGTTCCTTCACATG

     730     740     750     760     770     780
CGGGACATCGTGATTGCTGAAACCCTGGAGGACCTGGACAGAAACAAAGATGGCTATGTC

     790     800     810     820     830     840
CAGGTGGAGGAGTACATCGCGGATCTGTACTCAGCCGAGCCTGGGGAGGAGAGCCGGCG

     850     860     870     880     890     900
TGGGTGCAGACGGAGAGGCAGCAGTTCCGGGACTTCCGGGATCTGAACAAGGATGGGCAC

     910     920     930     940     950     960
CTGGATGGGAGTGAGGTGGGCCACTGGGTGCTGCCCCCTGCCCAGGACCAGCCCCCTGGTG

     970     980     990    1000    1010    1020
GAAGCCAACCACCTGCTGCACGAGAGCGACACGGACAAGGATGGGCGGCTGAGCAAAGCG

```



1030 1040 1050 1060 1070 1080  
 GAAATCCTGGGTAATTGGAACATGTTTGTGGGCAGTCAGGCCACCAACTATGGCGAGGAC  
 1090 1100 1110 1120 1130 1140  
 CTGACCCGGCACCACGATGAGCTGTGAGCACCGCGCACCTGCCACAGCCTCAGAGGCCCG  
 1150 1160 1170 1180 1190 1200  
 CACAATGACCGGAGGAGGGGCGCTGTGGTCTGGCCCCCTCCCTGTCCAGGCCCGCAGG  
 1210 1220 1230 1240 1250 1260  
 AGGCAGATGCAGTCCCAGGCATCCTCCTGCCCCCTGGGCTCTCAGGGACCCCTGGGTCCG  
 1270 1280 1290 1300 1310 1320  
 CTTCTGTCCCTGTACACCCCCAACCCAGGGAGGGGCTGTTCATAGTCCCAGAGGATAAG  
 1330 1340 1350 1360 1370 1380  
 CAATACCTATTTCTGACTGAGTCTCCAGCCCAGACCCAGGGACCCCTGGCCCCAAGCTC  
 1390 1400 1410 1420 1430 1440  
 AGCTCTAAGAACCGCCCCAACCCCTCCAGCTCCAAATCTGAGCCTCCACCACATAGACTG  
 1450 1460 1470 1480 1490 1500  
 AAATCCCCTGGCCCCAGCCCTCTCCTGCCTGGCCTGGGACACCTCCTCTCTGCC  
 1510 1520 1530 1540  
 AGGAGGCAATAAAAGCCAGCGCCGGGAAAAAAAAAAAAAAAAA

Seq ID NO: 38

Sequence of the predicted BMS37 polypeptide (Range: 1 to 328)

10 20 30 40 50 60  
 MMWRPSVLLLLLLLRHGAQKPSPDAGPHGQGRVHQAAPLSDAPHDDAHGNFYDHEAFL  
 70 80 90 100 110 120  
 GREVAKEFDQLTPEESQARLGRIVDRMDRAGDGDGWVSLAELRAWIAHTQQRHIRDSVSA  
 130 140 150 160 170 180  
 AWDTYDTRDGRVGEELRNATYGHYAPGEEFHDVEDAETYYKMLARDERRFRVADQDGD  
 190 200 210 220 230 240  
 SMATREELTAFLHPEEFPHMRDIVIAETLEDLDRNKDGYVQVEEYIADLYSAEPGEEPA  
 250 260 270 280 290 300  
 WVQTERQQFRDRLNKDGHLDGSEVGHVLPAPQDQPLVEANHLLHESDTDKDGRLSKA  
 310 320  
 EILGNWNMFVGSQATNYGEDLTRHDEL

Seq ID NO: 39

## Sequence of BMS42 cDNA (Range: 1 to 1990)

```

      10      20      30      40      50      60
CACGAGCCTGCCCCGGCCCCCGGCTCCAGCGAGCGAGCGGCGAGCAGGCGGCTCACAGAGG

      70      80      90     100     110     120
CCTGGCCGCCCCACGGAACCCGGGGCCCCGGCGCCGCCGCGATGTTTCCCCCGAGAA

     130     140     150     160     170     180
GACGTGGAACATCTCGTTCGCGGGCTGCGGCTTCCTCGGCGTCTACTACGTGCGCGTGGC

     190     200     210     220     230     240
CTCCTGCCTCCGCGAGCACGCGCCCTTCCTGGTGGCCAACGCCACGCACATCTACGGCGC

     250     260     270     280     290     300
CTCGGCCGGGGCGCTCACGGCCACGGCGCTGGTCACCGGGGTCTGCCTGGGTGAGGCTGG

     310     320     330     340     350     360
TGCCAAGTTCATTGAGGTATCTAAAGAGGCCCGGAAGCGGTTCCTGGGCCCCCTGCACCC

     370     380     390     400     410     420
CTCCTTGAACTGGTAAAGATCATCCGCAGTTTCCTGCTGAAGGTCTGCCTGCTGATAG

     430     440     450     460     470     480
CCATGAGCATGCCAGTGGGCGCCTGGGCATCTCCCTGACCCGCGTGTCTAGACGGCGAGAA

     490     500     510     520     530     540
TGTCATTATATCCCACTTCAACTCCAAGGACGAGCTCATCCAGGCCAATGTCTGCAGCGG

     550     560     570     580     590     600
TTTCATCCCCGTGTACTGTGGGCTCATCCCTCCCTCCCTCCAGGGGGTGCCTACGTGGA

     610     620     630     640     650     660
TGGTGGCATTTTCAGACAACCTGCCACTCTATGAGCTTAAGAACACCATCACAGTGTCCCC

     670     680     690     700     710     720
CTTCTCGGGCGAGAGTGACATCTGTCCGCAGGACAGCTCCACCAACATCCACGAGCTGCG

     730     740     750     760     770     780
GGTCACCAACACCAGCATCCAGTTCAACCTGCGCAACCTCTACCGCCTCTCCAAGGCCCT

     790     800     810     820     830     840
CTTCCCCCGGAGCCCCCTGGTGTGCGAGAGATGTGCAAGCAGGGATACCGGGATGGCCT

     850     860     870     880     890     900
GCGCTTTCTGCAGCGGAACGGCCTCCTGAACCGGCCCAACCCCTTGCTGGCGTTGCCCCC

     910     920     930     940     950     960
CGCCCGCCCCCACGGCCCAGAGGACAAGGACCAGGCAGTGGAGAGCGCCCAAGCGGAGGA

     970     980     990    1000    1010    1020
T TACTCGCAGCTGCCCCGAGAAGATCACATCCTGGAGCACCTGCCCCCGCGCTCAATGA

```

1030 1040 1050 1060 1070 1080  
GGCCCTGCTGGAGGCCTGCGTGGAGCCCACGGACCTGCTGACCACCTCTCCAACATGCT

1090 1100 1110 1120 1130 1140  
GCCTGTGCGTCTGGCCACGGCCATGATGGTGCCCTACACGCTGCCGCTGGAGAGCGCTCT

1150 1160 1170 1180 1190 1200  
GTCCCTCACCATCCGCTTGCTGGAGTGGCTGCCCGACGTTCCCGAGGACATCCGGTGGAT

1210 1220 1230 1240 1250 1260  
GAAGGAGCAGACGGGCAGCATCTGCCAGTACCTGGTGATGCGCGCCAAGAGGAAGCTGGG

1270 1280 1290 1300 1310 1320  
CAGGCACCTGCCCTCCAGGCTGCCGGAGCAGGTGGAGCTGCGCCGCGTCCAGTCCGCTGCC

1330 1340 1350 1360 1370 1380  
GTCCGTGCCGCTGTCTGCGCCGCTACAGAGAGGCACTGCCCGGCTGGATGCGCAACAA

1390 1400 1410 1420 1430 1440  
CCTCTCGCTGGGGGACGCGCTGGCCAAGTGGGAGGAGTGCCAGCGCCAGCTGCTGCTCGG

1450 1460 1470 1480 1490 1500  
CCTCTTCTGCACCAACGTGGCCTTCCCGCCGAAGCTCTGCGCATGCGCGCACCCGCCGA

1510 1520 1530 1540 1550 1560  
CCCGGCTCCCGCCCCCGGACCCAGCATCCCCGACGACCCAGCCGGCCGGCCTGCCCC

1570 1580 1590 1600 1610 1620  
CTTGCTGAGCACCCCTGCTCCCGAGGCCCCGCGCGTGATCGGGGCCCTGGGGCTGTGAGA

1630 1640 1650 1660 1670 1680  
CCCCGACCCCTCTCGAGGAACCCCTGCCTGAGACGCTCCATTACCACTGCGCAGTGAGATG

1690 1700 1710 1720 1730 1740  
AGGGGACTCACAGTTGCCAAGAGGGGTCTTTGCCGTGGGCCCCCTCGCCAGCCACTCACC

1750 1760 1770 1780 1790 1800  
AGCTGCACTGAGAGGGGAGGTTTCCACACCCCTCCCTGGGCGGCTGAGGCCCCGCGCAC

1810 1820 1830 1840 1850 1860  
CTGTGCCTTAATCTTCCCTCCCCTGTGCTGCCCAGCACCTCCCCCGCCCCCTTACTCCT

1870 1880 1890 1900 1910 1920  
GGGAACCTTTCAGCTGCCCTTCCCTCCCCGTTTTTCATGGCCTGCTGAAATATGTGTGTG

1930 1940 1950 1960 1970 1980  
AAGAATTATTTATTTTCGCCAAAGCACATGTAATAAATGCTGCAGCCCAGAAAAAAAAA

1990  
AAAAAAAAA

SEQ ID NO:40

Sequence of the predicted BMS42 p lypeptide (Range: 1 to 504)

10	20	30	40	50	60
<u>MFPREKTWNISFAGCGFLGVYVGVASCLREHAPFLVANATHLYGASAGALTATALVTGV</u>					
70	80	90	100	110	120
<u>CLGEAGAKFIEVSKEARKRFLGPLHPSFNLVKIIRSFLKVLPA</u> <u>DSHEHASGRLGISLTR</u>					
130	140	150	160	170	180
VSDGENVIISHFNSKDELIQANVCSGFIPVYCGLI					
PPSLQGVRYVDGGISDNLP					
LYELKN					
190	200	210	220	230	240
TITVSPFSGESD					
ICPQDSSTNIHEL					
RVTNTSIQFNLRNLYRLSKALFPPEPLV					
LREMCKQ					
250	260	270	280	290	300
GYRDGLRFLQRNGLLNRPNPL					
LALPPARPHGPEDKDQAVESAQAEDYSQLPGEDHILEHL					
310	320	330	340	350	360
PARLNEAL					
LEACVEPTDLLTTLSNMLPVRLATAMMVPYTLP					
LESALSFTIRLLEWLPDVP					
370	380	390	400	410	420
EDIRWMKEQTGSICQYLV					
MRKRKLGRHLPSRLPEQVELRRVQSLPSVPLSCAAYREALP					
430	440	450	460	470	480
GWMRNLSLGDALAKWEECQRQLLLGLFCTNVAF					
PPEALRMRAPADPAPAPADPASPOHQ					
490	500				
PAGPAPLLSTPAPEARPVIGALGL					

SEQ ID NO. 41

Sequence of BMS60 cDNA (Range: 1 to 684)

```

      10      20      30      40      50      60
ACCGTCATGCTCCAGTTCTTTGTGCACTTCCTGAGCCTTGTCTACCTGTACCGTGAGGCC

      70      80      90     100     110     120
CAGGCCCGGAGCCCCGAGAAGCAGGAGCAGTTCGTGGACTTGTACAAGGAGTTTGAGCCA

     130     140     150     160     170     180
AGCCTGGTCAACAGCACCGTCTACATCATGGCCATGGCCATGCAGATGGCCACCTTCGCC

     190     200     210     220     230     240
ATCAATTACAAAGGCCCGCCCTTCATGGAGAGCCTGCCCGAGAACAAGCCCTGGTGTGG

     250     260     270     280     290     300
AGTCTGGCAGTTTCACTCCTGGCCATCATTTGGCCTGCTCCTCGGCTCCTCGCCCGACTTC

     310     320     330     340     350     360
AACAGCCAGTTTGGCCTCGTGGACATCCCTGTGGAGTTCAAGCTGGTCATTGCCCAGGTC

     370     380     390     400     410     420
CTGCTCCTGGACTTCTGCCTGGCGCTCCTGGCCGACCGCGTCTGCAGTTCTTCTCTGGGG

     430     440     450     460     470     480
ACCCCGAAGCTGAAAGTGCCCTCCTGAGATGGCAGTGC'TGGTACCCACTGCCCACCCTGG

     490     500     510     520     530     540
CTGCCCGCTGGGCGGGAACCCCAACAGGGCCCCGGGAGGGAACCCCTGCCCCCAACCCCCA

     550     560     570     580     590     600
CAGCAAGGCTGTACAGTCTCGCCCTTGAAGACTGAGCTGGGACCCCCACAGCCATCCGC

     610     620     630     640     650     660
TGGCTTGGCCAGCAGAACCAGCCCCAAGCCAGCACCTTTGGTAAATAAAGCAGCATCTGA

     670     680
GATTTTAAAAAAAAAAAAAAAAAAAA

```

SEQ ID NO. 42

Sequence of the predicted BMS60 polypeptide (Range: 1 to 146)

```

      10      20      30      40      50      60
MLOFFVHFLSLVYLYREAQARSPEKQEQFVDLYKEFEPVLNSTVYIMAMAMQMATFAIN

      70      80      90     100     110     120
YKGPPFMESLPENKPLVWSLAVSLLAIIGLLLGSSPDFNSQFGLVDIPVEFKLVIAQVLL

     130     140
LDFCLALLADRVLQFFLGTPKLVPS

```

Seq ID No: 43

Sequence of BMS61 cDNA (Range: 1 to 1152)

```

      10      20      30      40      50      60
GGCAGAGGGCAGCCTCCCCCTCGCTCGCTCTCCTCTCTAGGGCCCCAGCGCAGCTC

      70      80      90     100     110     120
GGGAGCCCCGCGCACCAGGGCGCTAGGGGCACCGCGCACTAGAGGGACACCCGCCGCGCCT

      130     140     150     160     170     180
GGACAGCCCCCGCGGGCGCCCCCTCGCACCTCCTGCCCCGCGCGGGCCGCGCTCCCCCT

      190     200     210     220     230     240
CCCCCGCGCCTGTGTCCCCAGGGCGCAGGGCCGCGCGTCCAGCCCCAGACCCGCCGGGGT

      250     260     270     280     290     300
CCCTGGGGACGCGCCAGCCCCGGCAGTGGCTCGACGATGGAGGAGCCGCAGCGCGCCCGCT

      310     320     330     340     350     360
CGCACACAGTCACCACCACCGCCAGCTCCTTCGCAGAGAACTTCTCCACCAGCAGCAGCA

      370     380     390     400     410     420
GCTTCGCCTACGACCGGGAGTTCTTCGACCCCTGCCCGGCTTCTCATCGTGGCCGAGA

      430     440     450     460     470     480
TCGTTCTGGGGCTGCTGGTATGGACGCTTATTGCTGGAACGAGTACTTCCGGGTCCCCG

      490     500     510     520     530     540
CATTTGGCTGGGTCATGTTGTAGCTGTATTTTACTGGGTCTCACCCTCTTCTTCCTCA

      550     560     570     580     590     600
TTATCTACATAACAATGACCTACACCAGGATTCCCCAGGTGCCCTGGACAACAGTGGGCC

      610     620     630     640     650     660
TGTGCTTTAACGGCAGTGCCTTCGTCTTGTACCTCTCTGCCGCTGTTGTAGATGCATCTT

      670     680     690     700     710     720
CCGTCTCCCCTGAGAGGGACAGTCACAACCTTCAACAGCTGGGCGGCCTCATCGTTCTTTG

      730     740     750     760     770     780
CCTTCCTGGTCACCATCTGCTACGCTGGAAATACATATTTTCACTTTTATAGCATGGAGAT

      790     800     810     820     830     840
CCAGGACCATACAGTGATTTACCATTTTGATAATTAAAAGGAAAAAAGGAAGACTCT

      850     860     870     880     890     900
CACTGTAAAAACAGCTGTAGGTATAATGTATATTTCCAGAGAATGTATTTAACTAATTA

      910     920     930     940     950     960
ATGTTTTTTATATTCTTAAATTTGCTCACAAATTGTGGTTTGTACAATTAACTGGATA

      970     980     990    1000    1010    1020
CTTATTGCAAAGTGTTGTAGCTTATAATGAACTCTTAAGTATCTTATTAATGTATTAAT

```

1030 1040 1050 1060 1070 1080  
GTCTTCATAGATCATATTTTCTTAGACAATGTTTAAATAGATAAAATTGCTAATATTGAGA  
1090 1100 1110 1120 1130 1140  
ATGTGTCAAGTTTGTAAACCTAACTTTTAAGATGCCAGATTCTTTTGTGATTAAATGTTG  
1150  
CAAAATCCCAA

*Seq ID NO:44*

Sequence of the predicted BMS61 polypeptide (Range: 1 to 173)

10 20 30 40 50 60  
MEEPQRARSHTVTTTASSFAENFSTSSSSPAYDREFLRTLPGFLIVAEIVLGLLVWTLIA  
70 80 90 100 110 120  
GTEYFRVPAFGWVMFVAVFYWVLTVFFLIYYITMTYTRIPQVPWTTVGLCFNGSAFVLYL  
130 140 150 160 170  
SAAVVDASSVSPERDSHNFNNSWAASSFFAFLVTICYAGNTYFSFIAWRSRTIQ

SEQ ID NO:45 polyadenylation signal

AATAAA

SEQ ID NO:46 polyadenylation signal

ATTAAA





## INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

<b>(51) International Patent Classification 6 :</b> <b>C12N 15/12, C07K 14/47, 14/495, C12N 15/62, A61K 38/17, C07K 16/18, C12Q 1/68</b>	<b>A3</b>	<b>(11) International Publication Number:</b> <b>WO 99/33979</b> <b>(43) International Publication Date:</b> <b>8 July 1999 (08.07.99)</b>
<b>(21) International Application Number:</b> <b>PCT/US98/27008</b> <b>(22) International Filing Date:</b> <b>18 December 1998 (18.12.98)</b>  <b>(30) Priority Data:</b> 60/068,958      30 December 1997 (30.12.97)      US 60/101,603      24 September 1998 (24.09.98)      US 60/102,540      30 September 1998 (30.09.98)      US  <b>(71) Applicant:</b> <b>CHIRON CORPORATION [US/US]; 4560 Horton Street - R440, Emeryville, CA 94608 (US).</b>  <b>(72) Inventors:</b> <b>LIN, Haishan; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94622-8097 (US). CAO, Li; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94622-8097 (US).</b>  <b>(74) Agents:</b> <b>POTTER, Jane, E., R. et al.; Chiron Corporation, Intellectual Property - R440, P.O. Box 8097, Emeryville, CA 94662-8097 (US).</b>		<b>(81) Designated States:</b> <b>AU, CA, JP, European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE).</b>  <b>Published</b> <i>With international search report.</i> <i>Before the expiration of the time limit for amending the claims and to be republished in the event of the receipt of amendments.</i>  <b>(88) Date of publication of the international search report:</b> <b>16 September 1999 (16.09.99)</b>

**(54) Title:** **BONE MARROW SECRETED PROTEINS AND POLYNUCLEOTIDES**

**(57) Abstract**

Novel polynucleotides and secreted proteins encoded thereby are disclosed. The proteins can be used as therapeutics, for example, to stimulate blood cell generation in patients receiving cancer chemotherapy, to treat bone marrow transplantation patients, and to heal fractured bones. Polynucleotides of the invention can be used therapeutically, to provide proteins of the invention. Polynucleotides of the invention can also be used diagnostically, such as on polynucleotide arrays, to detect differential gene expression in diseased tissue compared with gene expression in normal tissue.

**FOR THE PURPOSES OF INFORMATION ONLY**

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece			TR	Turkey
BG	Bulgaria	HU	Hungary	ML	Mali	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MN	Mongolia	UA	Ukraine
BR	Brazil	IL	Israel	MR	Mauritania	UG	Uganda
BY	Belarus	IS	Iceland	MW	Malawi	US	United States of America
CA	Canada	IT	Italy	MX	Mexico	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NE	Niger	VN	Viet Nam
CG	Congo	KE	Kenya	NL	Netherlands	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NO	Norway	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	NZ	New Zealand		
CM	Cameroon	KR	Republic of Korea	PL	Poland		
CN	China	KZ	Kazakhstan	PT	Portugal		
CU	Cuba	LC	Saint Lucia	RO	Romania		
CZ	Czech Republic	LI	Liechtenstein	RU	Russian Federation		
DE	Germany	LK	Sri Lanka	SD	Sudan		
DK	Denmark	LR	Liberia	SE	Sweden		
EE	Estonia			SG	Singapore		

# INTERNATIONAL SEARCH REPORT

International Application No

PC., US 98/27008

## A. CLASSIFICATION OF SUBJECT MATTER

IPC 6 C12N15/12 C07K14/47 C07K14/495 C12N15/62 A61K38/17  
C07K16/18 C12Q1/68

According to International Patent Classification (IPC) or to both national classification and IPC

## B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

IPC 6 C07K C12N A61K C12Q

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practical, search terms used)

## C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
X	HWANG S-Y ET AL.: "Mus musculus cornichon mRNA (accession number AF022811)" EMBL SEQUENCE DATABASE, 3 October 1997 (1997-10-03), XP002099391 Heidelberg, Germany the whole document	1-3, 6-12, 14-18
Y	---	19-21
X	ROTH S ET AL.: "Cornichon and the EGF receptor signaling process are necessary for both anterior-posterior and dorsal-ventral pattern formation in Drosophila" CELL, vol. 81, 16 June 1995 (1995-06-16), pages 967-978, XP002099392 the whole document	12
	---	
	-/--	

☒ Further documents are listed in the continuation of box C.

☒ Patent family members are listed in annex.

### \* Special categories of cited documents :

- \*A\* document defining the general state of the art which is not considered to be of particular relevance
- \*E\* earlier document but published on or after the international filing date
- \*L\* document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)
- \*O\* document referring to an oral disclosure, use, exhibition or other means
- \*P\* document published prior to the international filing date but later than the priority date claimed

- \*T\* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
- \*X\* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
- \*Y\* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art.
- \*S\* document member of the same patent family

Date of the actual completion of the international search

12 April 1999

Date of mailing of the international search report

26. 07. 99

Name and mailing address of the ISA

European Patent Office, P.B. 5818 Patentlaan 2  
NL - 2280 HV Rijswijk  
Tel. (+31-70) 340-2040, Tx. 31 651 epo nl,  
Fax: (+31-70) 340-3016

Authorized officer

Oderwald, H

# INTERNATIONAL SEARCH REPORT

Int'l Application No  
PCT/US 98/27008

**C.(Continuation) DOCUMENTS CONSIDERED TO BE RELEVANT**

Category *	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
Y	LOCKHART D J ET AL: "EXPRESSION MONITORING BY HYBRIDIZATION TO HIGH-DENSITY OLIGONUCLEOTIDE ARRAYS" BIO/TECHNOLOGY, vol. 14, no. 13, December 1996 (1996-12), pages 1675-1680, XP002022521 the whole document	19-21
A	--- EP 0 409 472 A (CHIRON CORP) 23 January 1991 (1991-01-23) the whole document	1-21
A	--- WO 85 02863 A (BIOTECH AUSTRALIA PTY LTD ;UNIV AUSTRALIAN (AU)) 4 July 1985 (1985-07-04) the whole document	1-21
A	--- TASHIRO K ET AL: "SIGNAL SEQUENCE TRAP: A CLONING STRATEGY FOR SECRETED PROTEINS AND TYPE I MEMBRANE PROTEINS" SCIENCE, vol. 261, 30 July 1993 (1993-07-30), pages 600-603, XP000673204 the whole document	
T	--- PLISOV S Y ET AL.: "Homo sapiens cornichon mRNA (accession number AF104398)" EMBL SEQUENCE DATABASE, 29 December 1998 (1998-12-29), XP002099394 Heidelberg, Germany the whole document -----	1-3,6-18

# INTERNATIONAL SEARCH REPORT

International application No.

PCT/US 98/ 27008

## Box I Observations where certain claims were found unsearchable (Continuation of Item 1 of first sheet)

This International Search Report has not been established in respect of certain claims under Article 17(2)(a) for the following reasons:

1. ☐ Claims Nos.:  
because they relate to subject matter not required to be searched by this Authority, namely:
2. ☐ Claims Nos.:  
because they relate to parts of the International Application that do not comply with the prescribed requirements to such an extent that no meaningful International Search can be carried out, specifically:
3. ☐ Claims Nos.:  
because they are dependent claims and are not drafted in accordance with the second and third sentences of Rule 6.4(a).

## Box II Observations where unity of invention is lacking (Continuation of Item 2 of first sheet)

This International Searching Authority found multiple inventions in this international application, as follows:

see FURTHER INFORMATION sheet

1. ☐ As all required additional search fees were timely paid by the applicant, this International Search Report covers all searchable claims.
2. ☐ As all searchable claims could be searched without effort justifying an additional fee, this Authority did not invite payment of any additional fee.
3. ☐ As only some of the required additional search fees were timely paid by the applicant, this International Search Report covers only those claims for which fees were paid, specifically claims Nos.:
4. ☒ No required additional search fees were timely paid by the applicant. Consequently, this International Search Report is restricted to the invention first mentioned in the claims; it is covered by claims Nos.:

1-21 all partially (subject 1. on continuation sheet)

Remark on Protest

- ☐ The additional search fees were accompanied by the applicant's protest.
- ☐ No protest accompanied the payment of additional search fees.

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 98/27008

FURTHER INFORMATION CONTINUED FROM PCT/SA/ 210

1. Claims: 1-21 all partially

An isolated and purified polypeptide comprising SEQ ID NO: 2, a fragment thereof, a fusion protein comprising said polypeptides, an antibody binding to said polypeptides. An isolated and purified subgenomic polynucleotide encoding said polypeptides comprising SEQ ID NO:1, a fragment thereof, hybridizing polynucleotides, a construct comprising said polynucleotides, a host cell comprising said construct. A process for producing said polypeptides, a polynucleotide array comprising at least 12 nucleotides of said polynucleotide, a method of detecting differential gene expression comprising said polynucleotide array.

2. Claims: 1-21 all partially

same as invention 1 but comprising SEQ ID NO: 3 and 4.

3. Claims: 1-21 all partially

same as invention 1 but comprising SEQ ID NO: 5 and 6.

4. Claims: 1-21 all partially

same as invention 1 but comprising SEQ ID NO: 7 and 8.

5. Claims: 1-21 all partially

same as invention 1 but comprising SEQ ID NO: 9 and 10.

6. Claims: 1-21 all partially

same as invention 1 but comprising SEQ ID NO: 11 and 12.

7. Claims: 1-21 all partially

same as invention 1 but comprising SEQ ID NO: 13 and 14.

8. Claims: 1-21 all partially

same as invention 1 but comprising SEQ ID NO: 15 and 16.

9. Claims: 1-21 all partially

same as invention 1 but comprising SEQ ID NO: 17 and 18.

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 98/27008

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

10. Claims: 1-21 all partially  
same as invention 1 but comprising SEQ ID NO: 19 and 20.
11. Claims: 1-21 all partially  
same as invention 1 but comprising SEQ ID NO: 21 and 22.
12. Claims: 1-21 all partially  
same as invention 1 but comprising SEQ ID NO: 23 and 24.
13. Claims: 1-21 all partially  
same as invention 1 but comprising SEQ ID NO: 25 and 26.
14. Claims: 1-21 all partially  
same as invention 1 but comprising SEQ ID NO: 27 and 28.
15. Claims: 1-21 all partially  
same as invention 1 but comprising SEQ ID NO: 29 and 30.
16. Claims: 1-21 all partially  
same as invention 1 but comprising SEQ ID NO: 31 and 32.
17. Claims: 1-21 all partially  
same as invention 1 but comprising SEQ ID NO: 33 and 34.
18. Claims: 1-21 all partially  
same as invention 1 but comprising SEQ ID NO: 35 and 36.
19. Claims: 1-21 all partially  
same as invention 1 but comprising SEQ ID NO: 37 and 38.
20. Claims: 1-21 all partially  
same as invention 1 but comprising SEQ ID NO: 39 and 40.

# INTERNATIONAL SEARCH REPORT

International Application No. PCT/US 98/27008

FURTHER INFORMATION CONTINUED FROM PCT/ISA/ 210

21. Claims: 1-21 all partially

same as invention 1 but comprising SEQ ID NO: 41 and 42.

22. Claims: 1-21 all partially

same as invention 1 but comprising SEQ ID NO: 43 and 44.



# INTERNATIONAL SEARCH REPORT

Information on patent family members

International Application No

PCT/US 98/27008

Patent document cited in search report	Publication date	Patent family member(s)	Publication date
EP 0409472 A	23-01-1991	AT 114169 T	15-12-1994
		CA 2020729 A	20-01-1991
		DE 69014162 D	22-12-1994
		DE 69014162 T	11-05-1995
		DK 409472 T	16-01-1995
		ES 2063278 T	01-01-1995
		IE 66495 B	10-01-1996
		JP 3195495 A	27-08-1991
		PT 94732 A,B	20-03-1991
		US 5620867 A	15-04-1997
WO 8502863 A	04-07-1985	AT 85646 T	15-02-1993
		AU 570762 B	24-03-1988
		AU 3785485 A	12-07-1985
		DE 3486068 A	25-03-1993
		EP 0167548 A	15-01-1986

**THIS PAGE BLANK (USPTO)**